

- [1] Bhargava D: Phentolamine mesylate. *J Stomat Occ Med* 5, 53–56 (2012).
- [2] Daubländer M, Kämmerer PW, Willershausen B, Leckel M, Lauer HC, Buff S, Rösl B: Clinical use of epinephrine-reduced (1:400,000) articaine solution in short-time dental routine treatments: a multicenter study. *Clinical Oral Investigations* 16, 1289–1295 (2011).
- [3] Elmore S, Nusstein J, Drum M, Reader A, Beck M, Fowler S: Reversal of pulpal and soft tissue anesthesia by using phentolamine: a prospective randomized, single-blind study. *J Endod* 39, 429–434 (2013).
- [4] Heinzel F, Matthes K, Mechelke K, Nusser E: Die Kreislaufwirkung des Regitin beim gesunden Menschen. *Cardiologia* 21, 743–756 (1952).
- [5] Hersh EV, Moore PA, Papas AS, Goodson JM, Navalta LA, Rogy S, Rutherford B, Yagiela JA: Reversal of soft-tissue local anesthesia with phentolamine mesylate in adolescents and adults. *J Am Dent Assoc* 139, 1080–1093 (2008).
- [6] Hersh EV, Lindemeyer RG: Phentolamine mesylate for accelerating recovery from lip and tongue anesthesia. *Dent Clin North Am* 54, 631–642 (2010).
- [7] Kämmerer PW, Krämer N, Esch J, Pfau H, Uhlemann U, Piehlmeier L, Daubländer M: Epinephrine-reduced articaine solution (1:400,000) in paediatric dentistry: a multicentre non-interventional clinical trial. *European Archives of Paediatric Dentistry* 14, 89–95 (2013).
- [8] Kämmerer PW, Palarie V, Daubländer M, Bicer C, Shabazfar N, Brüllmann D, Al-Nawas B: Comparison of 4% articaine with epinephrine (1:100,000) and without epinephrine in inferior alveolar block for tooth extraction: double-blind randomized clinical trial of anesthetic efficacy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 113, 495–499 (2012).
- [9] Kerezoudis NP, Olgart L, Edwall L, Gazelius B, Nomikoa GG: Activation of sympathetic fibres in the pulp by electrical stimulation of rat incisor teeth. *Archs oral Biol* 37, 1013–1019 (1992).
- [10] Kerezoudis NP, Funato A, Edwall L, Olgart L: Activation of sympathetic nerves exerts an inhibitory influence on afferent nerve-induced vasodilation unrelated to vasoconstriction in rat dental pulp. *Acta Physiol Scand* 147, 27–35 (1993).
- [11] Koss MC: Differential neural activation of vascular α-adrenoceptors in oral tissues of cats. *European Journal of Pharmacology* 440, 53–59 (2002).
- [12] Laviola M, McGavin SK, Freer GA, Plancich G, Woodbury SC, Marinkovich S, Morrison R, Reader A, Rutherford RB, Yagiela JA: Randomized study of phentolamine mesylate for reversal of local anesthesia. *J Dent Res* 87, 635–639 (2008).
- [13] Maguire WM, Reisdorff EJ, Smith D, Wiegenstein JG: Epinephrine-induced vasospasm reversed by phentolamine digital block. *Am J Emerg Med* 8, 46–47 (1990).
- [14] McCleane GJ: Oral Phentolamine Mesylate in the treatment of complex regional pain syndrome. *The Ulster Medical Journal* 65, 87–88 (1996).
- [15] Moore PA, Hersh EV, Papas AS, Goodson JM, Yagiela JA, Rutherford B, Rogy S, Navalta L: Pharmacokinetics of lidocaine with epinephrine following local anesthesia reversal with phentolamine mesylate. *Anesth Prog* 55, 40–48 (2008).
- [16] Nishizawa S, Ichinohe T, Kaneko Y, Tissue blood flow reduction induced by remifentanil in rabbits and the effect of naloxone and phentolamin on these changes. *J Oral Maxillofac Surg* 70, 797–802 (2012).
- [17] Nourbakhsh N, Shirani F, Babaei M: Effect of phentolamine mesylate on duration of soft tissue local anesthesia in children. *Journal of Research in Pharmacy Practice* 1, 55–59 (2012).
- [18] Raja SN, Treede RD, Davis KD, Campbell JN: Systemic alpha-adrenergic blockade with phentolamine: a diagnostic test for sympathetically maintained pain. *Anesthesiology* 74, 691–698 (1991).
- [19] Rutherford B, Zeller JR, Thake D: Local and systemic toxicity of intraoral submucosal injections of phentolamine mesylate (OraVerse). *Anesth Prog* 56, 123–127 (2009)e.

- [20] Shir Y, Cameron LB, Raja SN, Bourke DL: The safety of intravenous phentolamine administration in patients with neuropathic pain. Anesth Analg 76, 1008–1011 (1993).
- [21] Stadler A, Daubländer M: Patienteneinschätzung der zahnärztlichen Lokalanästhesie-Befragungen zu Erfahrungen, zur Anamneseerhebung und Aufklärung. DZZ 59, 90–93 (2004).
- [22] Tavares M, Goodson JM, Studen-Pavlovich D, Yagiela JA, Navalta LA, Rogy S, Rutherford B, Gordon S, Papas AS: Reversal of soft-tissue local anesthesia with phentolamine mesylate in pediatric patients. J Am Dent Assoc 139, 1095–1104 (2008).
- [23] Yagiela JA: What's new with phentolamine mesylate: a reversal agent for local anaesthesia. SAAD Dig 27, 3–7 (2011).

Verzeichnis elektronischer Quellen

<http://www.fda.gov/downloads/Drugs/DevelopmentApprovalProcess/DevelopmentResources/ucm072805.pdf>

<http://www.fda.gov/downloads/Drugs/DevelopmentApprovalProcess/DevelopmentResources/ucm072807.pdf>

http://www.ineedce.com/courses/2088/PDF/1103cei_phentolamine_web.pdf