

Direkte adhäsive Versorgungen wurzelkanalbehandelter Zähne

- [1] Da Rosa Rodolfo PA, Donassollo TA, Cenci MS, Loguércio AD, Moraes RR, Bronkhorst EM, Opdam NJ, Demarco FF. 22 year clinical evaluation of the performance of two posterior composites with different filler characteristics. Dent Mater 2011; 27: 955–963.
- [2] Dammaschke T, Nykiel K, Sagheri D, Schäfer E. Influence of coronal restorations on the fracture resistance of root canal-treated premolar and molar teeth: a retrospective study. Aust Endod J 2013; 39: 48–56.
- [3] Deliperi S, Bardwell DN. Direct cuspal-coverage posterior resin composite restorations: A case report. Oper Dent 2006; 31: 143–150.
- [4] ElAyouti A, Serry MI, Geis-Gerstorfer J, Löst C. Influence of cusp coverage on the fracture resistance of premolars with endodontic access cavities. Int Endod J 2011; 44: 543–549.
- [5] Ernst CP. Die 20 beliebtesten Fehler beim Kleben. DFZ 2010; 11: 66–83.
- [6] Ernst CP. CEREC versus Straumann: Welche Teilkronen passt besser? Teil 1. ZMK 2013; 29: 184–193.
- [7] Ernst CP. Die Verwendung eines weiß-opaken Flow-Kompositen zur Markierung tiefer Kavitätenanteile in der Klasse II. Ästhetische Zahnmedizin 2004; 7: 160–169.
- [8] Ernst CP, Rullmann I, Janssen B, Willershausen B. Polymerization shrinkage stress of bulk fill resin composites. Vortrag auf der IADR Jahrestagung 2014.
<https://iadr.confex.com/iadr/13iags/webprogram/Paper170911.html>.
- [9] Ferracane J, Watts DC, Barghi N, Ernst CP, Rueggeberg FA, Shortall A, Price R, Strassler H. Der effiziente Einsatz von Lichtpolymerisationsgeräten – ein Leitfaden für Zahnärzte. ZMK 2014; 30: 166–180.
- [10] Ferrari M, Cagidiaco MC, Goracci C, Vichi A, Mason PN, Radovic I, Tay F. Long-term retrospective study of the clinical performance of fiber posts. Am J Dent 2007; 20:287–291.
- [11] Frankenberger R, Reinelt C, Roggendorf MJ, Lrämer N. Direkte oder indirekte Restaurationen im Seitenzahnbereich? Quintessenz 2010; 561: 575–579.
- [12] Goracci C, Grandini S, Bossù M, Bertelli E, Ferrari M. Laboratory assessment of the retentive potential of adhesive posts: a review. J Dent 2007; 35: 827–835.
- [13] Hickel R, Ernst CP, Haller B, Hugo B, Kunzelmann KH, Merte K, Ott K, Schmalz G, Staehle HJ. Direkte Kompositrestaurationen im Seitenzahnbereich – Indikation und Lebensdauer Zahnärztliche Mitteilungen 2005; 95: 74–76.
- [14] Juloski J, Fadda GM, Monticelli F, Fajó-Pascual M, Goracci C, Ferrari M. Four-year Survival of Endodontically Treated Premolars Restored with Fiber Posts. J Dent Res 2014 19; 93 (7 suppl): 52S–58S.
- [15] Kon M, Nicola Zitzmann NU, Weiger R, Krastl G. Postendodontische Restauration: eine Umfrage unter Zahnärzten der Schweiz. Schweiz Monatsschr Zahnmed 2013; 123: 1083–1088.
- [16] Manhart, J, Chen H, Hamm, G, Hickel R: Buonocore Memorial Lecture. Review of the clinical survival of direct and indirect restorations in posterior teeth of the permanent dentition. Oper Dent 2004;29: 481–508.
- [17] Mannocci F, Bertelli E, Sherriff M, Watson TF, Ford TR. Three-year clinical comparison of survival of endodontically treated teeth restored with either full cast coverage or with direct composite restoration. J Prosthet Dent 2002; 88:297–301.
- [18] Mohammadi N, Kahnamoii MA, Yeganeh PK, Navimipour EJ. Effect of fiber post and cusp coverage on fracture resistance of endodontically treated maxillary premolars directly restored with composite resin. J Endod. 2009; 35: 1428–1432.

- [19] [Mondelli RF](#), [Ishikirama SK](#), [de Oliveira Filho O](#), [Mondelli J](#). Fracture resistance of weakened teeth restored with condensable resin with and without cusp coverage. *J Appl Oral Sci* 2009;17: 161–165.
- [20] Nagasiri R, Chitmongkolsuk S: Long-term survival of endodontically treated molars without crown coverage: a retrospective cohort study. *J Prosthet Dent.* 2005; 93: 164–170.
- [21] [Naumann M](#), [Preuss A](#), [Rosentritt M](#). Effect of incomplete crown ferrules on load capacity of endodontically treated maxillary incisors restored with fiber posts, composite build-ups, and all-ceramic crowns: an in vitro evaluation after chewing simulation. *Acta Odontol Scand* 2006; 64: 31–36.
- [22] [Naumann M](#), [Koelpin M](#), [Beuer F](#), [Meyer-Lueckel H](#). 10-year survival evaluation for glass-fiber-supported postendodontic restoration: a prospective observational clinical study. *J Endod* 2012; 38: 432–435.
- [23] [Nothdurft FP](#), [Seidel E](#), [Gebhart F](#), [Naumann M](#), [Motter PJ](#), [Pospiech PR](#). Influence of endodontic posts on the fracture behavior of crowned premolars with Class II cavities. *J Dent* 2008; 36: 287–393.
- [24] Opdam NJM, Bronkhorst EM, Loomans BAC, Huysmans M-CDNM: 12-year survival of composite vs. amalgam restorations. *JDR Online First*, published on July 26, 2010.
- [25] Plotino G, Buono L, Grande NM, Lsamorgese V, Somma F. Fracture resistance of endodontically treated molars restored with extensive composite resin restorations. *J Prosthet Dent* 2008; 99: 225–232.
- [26] Rosentritt M, Behr M, Kolbeck C, Handel G. Flexural strength of restorative composites after different aging conditions. Vortrag auf der IADR Jahrestagung 2014. <https://iadr.confex.com/iadr/13iags/webprogram/Paper171330.html>
- [27] Rullmann I, Schattenberg A, Marx M, Willershausen B, Ernst CP: Spannungsoptische Messungen der Polymerisationsschrumpfungskraft schrumpfreduzierter Komposite. *Schweiz Monatsschr Zahnmed* 2012; 122: 8–12.
- [28] Scotti N, Rota R, Scanetti M, Paolono DS, Chisandussi G, Pasqualini D, Berutti E. Influence of adhesive techniques on fracture resistance of endodontically treated premolars with various residual wall thicknesses. *J Prosthet Dent* 2013; 110: 376382.
- [29] Staehle J. Restauration eines “hoffnungslosen Zahnes” mittels direkter Kompositkrone. *Quintessenz* 2006; 57: 531–537.
- [30] [Sterzenbach G](#), [Franke A](#), [Naumann M](#). Rigid versus flexible dentine-like endodontic posts-clinical testing of a biomechanical concept: seven-year results of a randomized controlled clinical pilot trial on endodontically treated abutment teeth with severe hard tissue loss. *J Endod* 2012; 38: 1557–1563.
- [31] [Xie KX](#), [Wang XY](#), [Gao XJ](#), [Yuan CY](#), [Li JX](#), [Chu CH](#). Fracture resistance of root filled premolar teeth restored with direct composite resin with or without cusp coverage. *Int Endod J.* 2012; 45: 524–529.