

- [1] Crossley K, Bennell K, Green S, Cowan S, McConnell J: Physical therapy for patellofemoral pain: a randomized, double-blinded, placebo-controlled trial. *Am J Sports Med* 30, 857–865 (2002).
- [2] Ferrario VF, Tartaglia GM, Luraghi FE, Sforza C: The use of surface electromyography as a tool in differentiating temporomandibular disorders from neck disorders. *Man Ther* 12, 372–379 (2007).
- [3] Glaros AG, Glass EG, Brockman D: Electromyographic data from TMD patients with myofascial pain and from matched control subjects: evidence for statistical, not clinical, significance. *J Orofac Pain* 11, 125–129 (1997).
- [4] Hellmann D, Brüstle F, Terebesi S, et al: Static balancing behaviour of the mandible. *Eur J Oral Sci* (2015); In Druck.
- [5] Hellmann D, Giannakopoulos NN, Blaser R, Eberhard L, Rues S, Schindler HJ: Long-term training effects on masticatory muscles. *J Oral Rehabil* 38, 912–920 (2011).
- [6] Hides JA, Jull GA, Richardson CA: Long-term effects of specific stabilizing exercises for first-episode low back pain. *Spine (Phila Pa 1976)* 26, E243–248 (2001).
- [7] Hides JA, Richardson CA, Jull GA: Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine (Phila Pa 1976)* 21, 2763–2769 (1996).
- [8] Hodges PW. Pain and motor control: From the laboratory to rehabilitation. *J Electromyogr Kinesiol* 21, 220–228 (2011).
- [9] Hodges PW, Richardson CA: Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine (Phila Pa 1976)* 21, 2640–2650 (1996).
- [10] Hodges PW, Tucker K: Moving differently in pain: a new theory to explain the adaptation to pain. *Pain* 152, S90–98 (2011).
- [11] Iida T, Komiyama O, Honki H, et al: Effect of a repeated jaw motor task on masseter muscle performance. *Arch Oral Biol* 60, 1625–1631 (2015).
- [12] Kumar A, Grigoriadis J, Trulsson M, Svensson P, Svensson KG: Effects of short-term training on behavioral learning and skill acquisition during intraoral fine motor task. *Neuroscience* 306, 10–17 (2015).
- [13] Lund J, Donga R, Widmer C, Stohler C: The pain-adaptation model: a discussion of the relationship between chronic musculoskeletal pain and motor activity. *Can J Physiol Pharmacol* 69, 683–694 (1991).
- [14] Magnusson T, Egermark I, Carlsson GE: A longitudinal epidemiologic study of signs and symptoms of temporomandibular disorders from 15 to 35 years of age. *J Orofac Pain* 14, 310–319 (2000).
- [15] McGill S: Low back disorders: evidence-based prevention and rehabilitation. *Human Kinetics* (2007).
- [16] Medicott MS, Harris SR: A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther* 86, 955–973 (2006).
- [17] Newell KM. Motor skill acquisition. *Annu Rev Psychol* 42, 213–237 (1991).

- [18] Pereira LJ, Steenks MH, de Wijer A, Speksnijder CM, van der Bilt A: Masticatory function in subacute TMD patients before and after treatment. *J Oral Rehabil* 36, 391–402 (2009).
- [19] Schindler HJ, Hugger A, Kordaß B, Türp JC: Grundlagen der Schienentherapie bei Myoarthropathien des Kausystems. *Z Kраниomandib Funkt* 6, 207–230 (2014).
- [20] Schindler HJ, Türp JC, Nilges P, Hugger A: Therapie bei Schmerzen der Kaumuskulatur: Aktualisierung der Empfehlungen. *Schmerz* 27, 243–252 (2013).
- [21] Sehm B, Taubert M, Conde V, et al: Structural brain plasticity in Parkinson's disease induced by balance training. *Neurobiol Aging* 35, 232–239 (2014).
- [22] Stolzenberg H: Bundes-Gesundheitssurvey (1998). Berlin: Robert-Koch-Institut (2000).
- [23] Stuge B, Laerum E, Kirkesola G, Vollestad N: The efficacy of a treatment program focusing on specific stabilizing exercises for pelvic girdle pain after pregnancy: a randomized controlled trial. *Spine (Phila Pa 1976)* 29, 351–359 (2004).
- [24] Svensson P: Masseter reflexes modulated by pain. *Mov Disord* 17 (Suppl 2), S45–48 (2002).
- [25] Tartaglia GM, Moreira Rodrigues da Silva MA, Bottini S, Sforza C, Ferrario VF: Masticatory muscle activity during maximum voluntary clench in different research diagnostic criteria for temporomandibular disorders (RDC/TMD) groups. *Man Ther* 13, 434–440 (2008).
- [26] Taubert M, Draganski B, Anwander A, et al: Dynamic properties of human brain structure: learning-related changes in cortical areas and associated fiber connections. *J Neurosci* 30, 11670–11677 (2010).
- [27] Türp JC, Schindler HJ, Pritsch M, Rong Q: Antero-posterior activity changes in the superficial masseter muscle after exposure to experimental pain. *Eur J Oral Sci* 110, 83–91 (2002).