

Erste Beschreibung einer Legionella-Infektion in Zusammenhang mit einer kontaminierten Dentaleinheit

- [1] Oppenheim, B. A., A. M. Sefton, O. N. Gill, J. E. Tyler, O. M. MC, J. M. Richards, P. J. Dennis, and T. G. Harrison, 1987, Widespread Legionella pneumophila contamination of dental stations in a dental school without apparent human infection: *Epidemiol Infect*, v. 99, no. 1, p. 159–166.
- [2] Lück, C., K. Neumann, B. Reitemeier, and L. Jatzwauk, 2011, Legionellen in Dentaleinheiten – Infektionsrisiko für Patienten und Personal?: *Zahnmedizin up2date*, v. 5, p. 347–361
- [3] Atlas, R. M., J. F. Williams, and M. K. Huntington, 1995, Legionella contamination of dental-unit waters: *Appl Environ Microbiol*, v. 61, no. 4, p. 1208–1213.
- [4] Pankhurst, C. L., W. Coulter, J. J. Philpott-Howard, T. Harrison, F. Warburton, S. Platt, S. Surman, and S. Challacombe, 2003, Prevalence of legionella waterline contamination and Legionella pneumophila antibodies in general dental practitioners in London and rural Northern Ireland: *Br Dent J*, v. 195, no. 10, p. 591–594.
- [5] Singh, T., and M. M. Coogan, 2005, Isolation of pathogenic Legionella species and legionella-laden amoebae in dental unit waterlines: *J Hosp Infect*, v. 61, no. 3, p. 257–262.
- [6] Ma'ayeh, S. Y., A. S. Al-Hiyasat, M. Y. Hindiyeh, and Y. S. Khader, 2008, Legionella pneumophila contamination of a dental unit water line system in a dental teaching centre: *Int J Dent Hyg*, v. 6, no. 1, p. 48–55.
- [7] Ricci, M. L., S. Fontana, F. Pinci, E. Fiumana, M. F. Pedna, P. Farolfi, M. A. B. Sabattini, and M. Scaturro, 2012, Pneumonia associated with a dental unit waterline: *The Lancet*, v. 379, no. 9816, p. 684.
- [8] Helbig, J. H., P. C. Lück, Y. A. Knirel, W. Witzleb, and U. Zähringer, 1995, Molecular characterization of a virulence-associated epitope on the lipopolysaccharide of Legionella pneumophila serogroup 1: *Epidemiol Infect*, v. 115, no. 1, p. 71–78.
- [9] Borchardt, J., J. H. Helbig, and P. C. Lück, 2008, Occurrence and distribution of sequence types among Legionella pneumophila strains isolated from patients in Germany: common features and differences to other regions of the world: *Eur J Clin Microbiol Infect Dis*, v. 27, no. 1, p. 29–36.
- [10] Harrison, T. G., B. Afshar, N. Doshi, N. K. Fry, and J. V. Lee, 2009, Distribution of Legionella pneumophila serogroups, monoclonal antibody subgroups and DNA sequence types in recent clinical and environmental isolates from England and Wales (2000–2008): *Eur J Clin Microbiol Infect Dis*, v. 28, no. 7, p. 781–791.
- [11] Franco, F. S., D. Spratt, J. C. Leao and S. R. Porter, 2005, Biofilm formation and control in dental unit waterlines. *Biofilms* (2005) 2, C_ Cambridge University Press, p. 9–17
- [12] Cirillo, J. D., S. Falkow, and L. S. Tompkins, 1994, Growth of Legionella pneumophila in Acanthamoeba castellanii enhances invasion: *Infect Immun*, v. 62, no. 8, p. 3254–3261.
- [13] O'Brien, J. Sarah, and M. Bophal, 1993, Legionnaires disease: The infective dose paradox: *Lancet*, v. 342, no. 8862, p. 8862.
- [14] Lück, P. C., C. Windmeier, T. Schreiber, I. Steinfeld, L. Gerber, and K. Teuber, 2010, Fallbericht: Reiseassoziierte Legionella-Pneumonie nach Aufenthalt in einem deutschen Hotel: *Epid.Bulletin*, no. 41, p. 405–407.
- [15] Robert-Koch-Institut, 2006, Infektionsprävention in der Zahnheilkunde – Anforderungen an die Hygiene: *Bundes Ges.-Blatt*, v. 49, p. 375–394.