

Kieferorthopädische Extraktionstherapie bei Molaren-Inzisiven-Hypomineralisation (MIH)

- [1] Weerheijm KL, Jalevik B, Alaluusua S: Molar-incisor hypomineralisation. *Caries Res* 35, 390–391 (2001).
- [2] Weerheijm KL, Duggal M, Mejare I et al.: Judgement criteria for molar incisor hypomineralisation (MIH) in epidemiologic studies: a summary of the European meeting on MIH held in Athens, 2003. *Eur J Paediatr Dent* 4, 110–113 (2003).
- [3] Ogden AR, Pinhasi R, White WJ: Nothing new under the heavens: MIH in the past? *Eur Arch Paediatr Dent* 9, 166–171 (2008).
- [4] Knapp V, Nies S: Molar-Incisor-Hypomineralization. *Zahnmedizin up2date* 3, 491–510 (2009).
- [5] Ostertag S: Die Molaren-Inzisiven-Hypomineralisation – ein bedeutendes Problem in der Kinderzahnmedizin. *Quintessenz* 62, 1553–1557 (2011).
- [6] Jalevik B: Prevalence and diagnosis of molar-incisor-hypomineralisation (MIH): A systematic review. *Eur J Paediatr Dent* 11, 59–64 (2010).
- [7] Heitmüller D, Hickel R, Kühnisch J: Diagnostik der Molaren-Inzisior-Hypomineralisation. *Quintessenz* 62, 1559–1563 (2011).
- [8] Kellerhof N: Ätiologie der Molaren-Inzisiven-Hypomineralisation. *Quintessenz* 62, 1565–1571 (2011).
- [9] Jalevik B, Dietz W, Noren JG: Scanning electron micrograph analysis of hypomineralized enamel in permanent first molars. *Int J Paediatr Dent* 15, 233–240 (2005).
- [10] Lygidakis NA, Wong F, Jalevik B et al.: Best clinical practice guidance for clinicians dealing with children presenting with molar-incisor-hypomineralisation (MIH): An EAPD Policy Document. *Eur Arch Paediatr Dent* 11, 75–81 (2010).
- [11] Jalevik B, Klingberg GA: (2002) Dental treatment, dental fear and behaviour management problems in children with severe enamel hypomineralization of their permanent first molars. *Int J Paediatr Dent* 12, 24–32 (2002).
- [12] Schätzle M, Patcas R: Idealer Extraktionszeitpunkt der ersten bleibenden Molaren – eine Literaturübersicht. *Quintessenz* 62, 1631–1635 (2011).
- [13] Eichenberger M, Erb J, Zwahlen M et al.: The timing of extraction of non-restorable first permanent molars: a systematic review. *Eur J Paediatr Dent* 16, 272–278 (2015).
- [14] Williams JK, Gowans AJ: Hypomineralised first permanent molars and the orthodontist. *Eur J Paediatr Dent* 4, 129–132 (2003).
- [15] Fayle SA: Molar incisor hypomineralisation: restorative management. *Eur J Paediatr Dent* 4, 121–126 (2003).
- [16] Steffen R: Therapie der Molaren-Inzisiven-Hypomineralisation in einem schwierigen Umfeld: Praktische Konzepte für den Behandlungsalltag. *Quintessenz* 62, 1613–1623 (2011).
- [17] Lygidakis NA: Treatment modalities in children with teeth affected by molar-incisor enamel hypomineralisation (MIH): A systematic review. *Eur Arch Paediatr Dent* 11, 65–74 (2010).
- [18] Onat H, Tosun G: Molar incisor hypomineralization. *J Paediatr Dent* 1, 53 (2013).
- [19] Feierabend S: Spätfolgen der Molaren-Inzisiven-Hypomineralisation: Möglichkeiten der prothetischen Langzeittherapie. *Quintessenz* 62, 1637–1643 (2011).
- [20] Cobourne MT, Williams A, Harrison M: National clinical guidelines for the extraction of first permanent molars in children. *Brit Dent J* 217, 643–648 (2014).
- [21] Koch MJ, Garcia-Godoy F: The clinical performance of laboratory-fabricated crowns placed on first permanent molars with developmental defects. *J Am Dent Assoc* 131, 1285–1290 (2000).

- [22] Zagdwon AM, Fayle SA, Pollard MA: A prospective clinical trial comparing preformed metal crowns and cast restorations for defective first permanent molars. *Eur J Paediatr Dent* 4, 138–142 (2003).
- [23] Jalevik B, Moller M: Evaluation of spontaneous space closure and development of permanent dentition after extraction of hypomineralized permanent first molars. *Int J Paediatr Dent* 17, 328–335 (2007).
- [24] Breunig A, Kirschneck C: Kieferorthopädische Therapie von Nichtanlagen – Lückenschluss oder Lückenöffnung? *Quintessenz* 67, 1–11 (2016).
- [25] Gill DS, Lee RT, Tredwin CJ: Treatment planning for the loss of first permanent molars. *Dent Update* 28, 304–308 (2001).
- [26] Sandler PJ, Atkinson R, Murray AM: For four sixes. *Am J Orthod Dentofacial Orthop* 117, 418–434 (2000).
- [27] Schopf P: Curriculum Kieferorthopädie: Werkstoffe. Festsitzende Apparaturen. Kieferorthopädische Therapie. Interdisziplinäre Aspekte. Anhang: Kieferorthopädische Abrechnung, 4., überarb. und erw. Aufl. Quintessenz-Verlag, Berlin (2008).
- [28] Komposch G: Extraction of 1st permanent molars within the framework of orthodontic treatment. Indications, timing and clinical problems (Die Sechsjahrmolarenextraktion im Rahmen der kieferorthopädischen Behandlung. Indikation, Zeitpunkt und klinische Problematik). *Dtsch Zahnärztl Z* 41, 100–104 (1986).
- [29] Thilander B, Skagius S: Orthodontic sequelae of extraction of permanent first molars. A longitudinal study. *Rep Congr Eur Orthod Soc*, 429–442 (1970).
- [30] Ong DC-V, Bleakley JE: Compromised first permanent molars: an orthodontic perspective. *Aust Dent J* 55, 2–14; quiz 105 (2010).
- [31] Baume LJ: Auswirkungen der Extraktion von Zähnen auf das deforme Gebiß. *Schweiz Mschr Zhlk* 49/50, 295–337/45–64 (1939/1940).
- [32] Ay S, Agar U, Bicakci AA et al.: Changes in mandibular third molar angle and position after unilateral mandibular first molar extraction. *Am J Orthod Dentofacial Orthop* 129, 36–41 (2006).
- [33] Livas C, Halazonetis DJ, Booij JW et al.: Extraction of maxillary first molars improves second and third molar inclinations in Class II Division 1 malocclusion. *Am J Orthod Dentofacial Orthop* 140, 377–382 (2011).
- [34] Bayram M, Ozer M, Arici S: Effects of first molar extraction on third molar angulation and eruption space. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 107, e14–20 (2009).
- [35] Mejare I, Bergman E, Grindeffjord M: Hypomineralized molars and incisors of unknown origin: treatment outcome at age 18 years. *Int J Paediatr Dent* 15, 20–28 (2005).
- [36] William V, Messer LB, Burrow MF: Molar incisor hypomineralization: review and recommendations for clinical management. *Pediatr Dent* 28, 224–232 (2006).
- [37] Jacobs C, Jacobs-Muller C, Luley C et al.: Orthodontic space closure after first molar extraction without skeletal anchorage. *J Orofac Orthop* 72, 51–60 (2011).