

**Bewertung des Maskierungseffekts durch Kariesinfiltration bei post-orthodontischen, kariösen Initialläsionen: 1-Jahres-Follow-up**

- [1] Sundararaj D, et al. Critical evaluation of incidence and prevalence of white spot lesions during fixed orthodontic appliance treatment: A meta-analysis. *J Int Soc Prev Community Dent.* 2015; 5 (6): 433-9.
- [2] Boersma JG, et al. Caries prevalence measured with QLF after treatment with fixed orthodontic appliances: influencing factors. *Caries Res.* 2005; 39 (1): 41-7.
- [3] Ogaard B. Prevalence of white spot lesions in 19-year-olds: a study on untreated and orthodontically treated persons 5 years after treatment. *Am J Orthod Dentofacial Orthop.* 1989; 96 (5): 423-7.
- [4] Kamber R, et al. Efficacy of sealants and bonding materials during fixed orthodontic treatment to prevent enamel demineralization: a systematic review and meta-analysis. *Sci Rep.* 2021; 11 (1): 16556.
- [5] Knosel M, et al. External bleaching effect on the color and luminosity of inactive white-spot lesions after fixed orthodontic appliances. *Angle Orthod.* 2007; 77 (4): 646-52.
- [6] Bourouni S, et al. Efficacy of resin infiltration to mask post-orthodontic or non-post-orthodontic white spot lesions or fluorosis - a systematic review and meta-analysis. *Clin Oral Investig.* 2021; 25(8): 4711-4719.
- [7] Kannan A, Padmanabhan S. Comparative evaluation of Icon(R) resin infiltration and Clinpro XT varnish on colour and fluorescence changes of white spot lesions: a randomized controlled trial. *Prog Orthod.* 2019; 20 (1): 23.
- [8] Willmot DR. White lesions after orthodontic treatment: does low fluoride make a difference? *J Orthod.* 2004; 31(3): 235-42; discussion 202.
- [9] Schmidlin PR, et al. Histological, morphological, profilometric and optical changes of human tooth enamel after microabrasion. *Am J Dent.* 2003; 16 Spec No: 4A-8A.
- [10] Knosel M, Eckstein A, Helms HJ. Durability of esthetic improvement following Icon resin infiltration of multibracket-induced white spot lesions compared with no therapy over 6 months: a single-center, split-mouth, randomized clinical trial. *Am J Orthod Dentofacial Orthop.* 2013; 144 (1): 86-96.
- [11] Kobbe C, et al. Evaluation of the value of re-wetting prior to resin infiltration of post-orthodontic caries lesions. *J Dent.* 2019; 91: 103243.
- [12] Jansen EE, et al. Do bleaching gels affect the stability of the masking and caries-arresting effects of caries infiltration-in vitro. *Clin Oral Investig.* 2021; 25 (6): 4011-4021.
- [13] Ceci M, et al. Resin infiltrant for non-cavitated caries lesions: evaluation of color stability. *J Clin Exp Dent.* 2017; 9 (2): e231-e237.
- [14] Paris S, et al. Masking of white spot lesions by resin infiltration in vitro. *J Dent.* 2013; 41 Suppl 5: e28-34.
- [15] Gu X, et al. Esthetic improvements of postorthodontic white-spot lesions treated with resin infiltration and microabrasion: A split-mouth, randomized clinical trial. *Angle Orthod.* 2019; 89 (3): 372-377.
- [16] Knosel M, Eckstein A, Helms HJ. Long-term follow-up of camouflage effects following resin infiltration of post orthodontic white-spot lesions in vivo. *Angle Orthod.* 2019; 89 (1): 33-39.
- [17] Knaup I, et al. Correlation of quantitative light-induced fluorescence and qualitative visual rating in infiltrated post-orthodontic white spot lesions. *Eur J Orthod.* 2022.

- [18] Wierichs RJ, Kobbe C, Meyer-Lueckel H. Wie der Maskierungserfolg von postorthodontischen White-Spot-Läsionen vor der Infiltration abgeschätzt werden kann. Dtsch Zahnärztl Z. 2020; 75(5): 60-66.
- [19] Mazur M, et al. Objective and subjective aesthetic performance of icon(R) treatment for enamel hypomineralization lesions in young adolescents: A retrospective single center study. J Dent. 2018; 68: 104-108.
- [20] Andrade R, et al. Clinical evaluation of the immediate masking effect of enamel white spot lesions treated with an infiltrant resin. Int J Esthet Dent. 2020; 15 (3): 306-316.
- [21] Kim S, et al. The evaluation of resin infiltration for masking labial enamel white spot lesions. Int J Paediatr Dent. 2011; 21 (4): 241-8.
- [22] Holmen L, Thylstrup A, Artun J. Surface changes during the arrest of active enamel carious lesions in vivo. A scanning electron microscope study. Acta Odontol Scand. 1987; 45 (6): 383-90.
- [23] Wierichs RJ, et al. Evaluation of the masking efficacy of caries infiltration in post-orthodontic initial caries lesions: 1-year follow-up. Clin Oral Investig. 2023.