ZMK 10/2022 (38) 540-543

Dr. Walter Denner

Die suffiziente Behandlung angrenzender approximaler Defekte

[1] DGZ, DGZMK. Kompositrestaurationen im Seitenzahnbereich; S1-Handlungsempfehlung (Langversion). AWMF-Registernummer: 083-028. Stand: 2016 Oktober
[2] Ausiello P, Ciaramella S, Garcia-Godoy F, Martorelli M, Sorrentino R, Gloria A. Stress distribution of bulk-fill resin composite in class II restorations. Am J Dentistry. 2017; 30: 227-232.
[3] Heintze SD, Rousson V. Clinical effectiveness of direct class II restorations - a meta-analysis. J Adhes Dent. 2012; 14: 407-431.
[4] Hugo B. Oszillierende Verfahren in der Präparationstechnik (Teil 1). Swiss Dent J. 1999; 109: 140-153.
[5] Denner W. Eine minimalinvasive Technik zur einfachen Restauration approximaler Kavitäten. Hole-in-one-Technik. Die Quintessenz. 2016; 67: 583-589.
[6] Putignano A, Cerutti A, Mangani F. Adhäsivtechnik in der ästhetischen Zahnheilkunde. Berlin: Quintessence Publishing Deutschland. 2009.
[7] Braga RR, Ballester RY, Ferracane JL. Factors involved in the development of polymerization shrinkage stress in resin-composites: a systematic review. Dental materials: official publication of the Academy of Dental Materials. 2005; 21: 962-970.
[8] Bichacho N. The centripetal build-up for composite resin posterior restorations. Pract Periodontics Aesthet Dent. 1994; 6: 17-23; quiz 24.