

ZMK 6/2023, 270-274

Prof. Dr. Claus-Peter Ernst

Das Inlay als Problemlöser bei Aufbissbeschwerden?

[1] Abbott P, Leow N. Predictable management of cracked teeth with reversible pulpitis. *Aust Dent J.* 2009; 54: 306-315.

[2] Abrams S: Improving the way to detect cracks in teeth. *Dent Today* 2013; 32: 104, 106.

[3] Aleixo AR, Guiraldo RD, Fugolin AP, Berger SB, Consani RL, Correr AB, Gonini-Júnior A, Lopes MB. Evaluation of contraction stress, conversion degree, and cross-link density in low-shrinkage composites. *Photomed Laser Surg.* 2014; 32: 267-273.

[4] Ardu S, Duc O, Di Bella E, Krejci I. Color stability of recent composite resins. *Odontology* 2017; 105: 29-35.

[5] Ardu S, Duc O, Di Bella E, Krejci I, Daher R. Color stability of different composite resins after polishing. *Odontology* 2018; 106: 328-333.

[6] Banerji S, Mehta SB, Millar BJ: Cracked tooth syndrome. Part 1: aetiology and diagnosis. *Br Dent J.* 2010 22; 208: 459–463.

[7] Banerji S, Mehta SB, Millar BJ. Cracked tooth syndrome. Part 2: restorative options for the management of cracked tooth syndrome. *Br Dent J.* 2010; 208: 503-514.

[8] Barreto BC, Van Meerbeek B, Van Ende A, Sousa SJ, Silva GR, Soares PV, Soares CJ. Biomechanical Behavior of Extensively Restored Premolars: Cusp Deformation, Marginal Integrity, and Fracture Resistance. *J Adhes Dent.* 2015; 17: 213-218.

[9] Blunck U, Ilie N. Lichtpolymerisation heute. *zm* 2019; 109: 2666-2676.

[10] Boaro LC, Gonçalves F, Guimarães TC, Ferracane JL, Versluis A, Braga RR. Polymerization stress, shrinkage and elastic modulus of current low-shrinkage restorative composites. *Dent Mater.* 2010; 26: 1144-1150.

[11] Boaro LC, Fróes-Salgado NR, Gajewski VE, Bicalho AA, Valdivia AD, Soares CJ, Miranda Júnior WG, Braga RR. Correlation between polymerization stress and interfacial integrity of composites restorations assessed by different in vitro tests. *Dent Mater.* 2014; 30: 984-992.

[12] Christensen GJ. Clinicians Report, A publication of CR Foundation, Provo, UT, USA, Newsletter April 2014, Volume 7, Issue 4.

[13] de Oliveira DC, Rovaris K, Hass V, Souza-Júnior EJ, Haiter-Neto F, Sinhoreti MA. Effect of low shrinkage monomers on physicochemical properties of dental resin composites. *Braz Dent J.* 2015; 26: 272-276.

[14] Ehrmann EH, Tyas MJ: Cracked tooth syndrome. *Aust Dent J.* 1990; 35: 390–391.

[15] Ernst CP. Die 20 beliebtesten Fehler beim Kleben. *Der Freie Zahnarzt* 2010; 11: 66-82.

[16] Ernst CP. Lichtpolymerisation. In: Frankenberger, R.: *Adhäsive Zahnheilkunde*, Deutscher Zahnärzte-Verlag, Köln, 2013: 35-80.

[17] Ernst CP. Vom Gold-Standard zur Gold-Alternative. *ZMK* 2015; 31, 214-226.

- [18] Ernst CP. Keramikteilkronen als Versorgungsoption von Zähnen mit Infraktionen. All ceramic partial crowns as a treatment option for cracked, but symptomless teeth. Deutsche Zahnärztl Z. 2015; 70 (3):165-173.
- [19] Ernst CP, Price RB, Callaway A, Massek A, Schwarm H, Rullmann I, Willershausen B, Ehlers V. Visible Light Curing Devices - Irradiance and Use in 302 German Dental Offices. J Adhes Dent. 2018; 20: 41-55.
- [20] Ernst CP. Augen auf beim Lampenkauf. ZMK 2019; 35; 14-20.
- [21] Ernst CP. Ein einziges Komposit – eine Farbe für alles? ZMK 2019; 35: 783-785.
- [22] Ernst CP. Komposit-Restaurationen: Differenzierte Materialauswahl bei benachbarten Kavitäten. ZMK 2020; 36: 599-603.
- [23] Ernst CP. Neu: etabliertes Komposit vereint universelle Farblösung mit sehr guten mechanischen Eigenschaften. ZMK 2020; 36: 776-780.
- [24] Ernst CP. Direkte Kompositrestaurationen als valide Alternative bei Inlay-Neuversorgungen. ZMK 2021; 37: 384-390.
- [25] Ernst CP. Omnichroma Flow- die perfekte Ergänzung zum pastösen Original. ZMK 2022; 38: 176-179.
- [26] Ernst CP. Die nächste Generation Bulk-Flowables: Fokus Ästhetik. ZMK 2022; 38 446-459.
- [27] Ferracane F, Watts DC, Barghi N, Ernst CP, Rueggeberg FA, Shortall A, Price R. Der effiziente Einsatz von Lichtpolymerisationsgeräten – ein Leitfaden für Zahnärzte. ZMK 2014; 30 (4): 166-180.
- [28] Ferracane JL, Hilton TJ. Polymerization stress - Is it clinically meaningful? Dent Mater 2016; 32: 1-10.
- [29] Frankenberger R. Composite first – Ceramic second: 25 Jahre Erfahrungen mit Komposit im Front- und Seitenzahnbereich. Quintessenz 2018; 69 (5): 490–497.
- [30] Frankenberger R, Dudek MC, Krämer N, Winter J, Roggendorf MJ. Die 10 beliebtesten Fehler in der Adhäsivtechnik Deutsche Zahnärztl Z. 2022; 77 (4):238-244.
- [31] Geurtsen W. The cracked-tooth syndrome: clinical features and case reports. Int J Periodontics Restorative Dent. 1992; 12: 395–405.
- [32] Geurtsen W, García-Godoy F: Bonded restorations for the prevention and treatment of the cracked-tooth syndrome. Am J Dent .1999; 12: 266–270.
- [33] Geurtsen W, Schwarze T, Günay H: Diagnosis, therapy, and prevention of the cracked tooth syndrome. Quintessence Int. 2003; 34: 409–417.
- [34] Heintze SD, Loguercio AD, Hanzen TA, Reis A, Rousson V. Clinical efficacy of resin-based direct posterior restorations and glass-ionomer restorations - An updated meta-analysis of clinical outcome parameters. Dent Mater. 2022; 38 (5): e109-e135.
- [35] Jacker-Guhr S: Infraktionen. Dtsch Zahnärztl Z. 2014; 69: 429–431.
- [36] Kahler W. The cracked tooth conundrum: terminology, classification, diagnosis, and management. Am J Dent. 2008; 21: 275-282.
- [37] Kim SY, Kim SH, Cho SB, Lee GO, Yang SE. Different treatment protocols for different pulpal and periapical diagnoses of 72 cracked teeth. J Endod. 2013; 39: 449-452.

- [38] Koottathape N, Takahashi H, Iwasaki N, Kanehira M, Finger WJ. Quantitative wear and wear damage analysis of composite resins in vitro. *J Mech Behav Biomed Mater.* 2014; 29:508-516.
- [39] Koplin, da Silva Rodrigues G, Jaeger R. Calculating internal stress during curing of dental composites. *J Dent Res.* 2009; 88 (Spec Iss B): 145.
- [40] Lubisich EB, Hilton TJ, Ferracane J. Cracked teeth: a review of the literature. *J Esthet Restor Dent.* 2010; 22: 158–167.
- [41] Lynch CD, Opdam NJ, Hickel R, Brunton PA, Gurgan S, Kakaboura A, Shearer AC, Vanherle G, Wilson NH. Guidance on posterior resin composites: Academy of Operative Dentistry - European Section. *J Dent.* 2014; 42: 377-383.
- [42] Magno MB, Rodrigues Nascimento GC, da Rocha YS, Ribeiro BD, Loretto SC, Maia LC. Silorane-based composite resin restorations are not better than conventional composites – a meta-analysis of clinical studies. *J Adhes Dent* 2016; 18: 375-386.
- [43] Marchesi G, Breschi L, Antonioli F, Di Lenarda R, Ferracane J, Cadenaro M. Contraction stress of low-shrinkage composite materials assessed with different testing systems. *Dent Mater.* 2010; 26: 947-953.
- [44] Pallesen U, van Dijken JW, Halken J, Hallonsten AL, Höigaard R. A prospective 8-year follow-up of posterior resin composite restorations in permanent teeth of children and adolescents in Public Dental Health Service: reasons for replacement. *Clin Oral Investig.* 2014; 18: 819-827.
- [45] Rullmann I, Schattenberg A, Marx M, Willershausen B, Ernst C-P. Photo-elastic determination of polymerization shrinkage stress in low-shrinkage resin composites. *Schweiz Monatsschr Zahnmed.* 2012; 22: 294-299.
- [46] Opdam NJ, Roeters JM. The effectiveness of bonded composite restorations in the treatment of painful, cracked teeth: six-month clinical evaluation. *Oper Dent.* 2003; 28: 327-333.
- [47] Opdam NJ, Roeters JJ, Loomans BA, Bronkhorst EM. Seven-year clinical evaluation of painful cracked teeth restored with a direct composite restoration. *J Endod.* 2008; 34: 808-811.
- [48] Peumans M, Politano G, Van Meerbeek B. Effective Protocol for Daily High-quality Direct Posterior Composite Restorations. Cavity Preparation and Design. *J Adhes Dent.* 2020; 22: 581 -596.
- [49] Peumans M, Venuti P, Politano G, Van Meerbeek B. Effective Protocol for Daily High-quality Direct Posterior Composite Restorations. The Interdental Anatomy of the Class-2 Composite Restoration. *J Adhes Dent.* 2021; 23: 21-34.
- [50] Schmidt C, Illie N. The mechanical stability of nano-hybrid composites with new methacrylate monomers for matrix compositions. *Dent Mater.* 2012; 28: 152-159.
- [51] Seo DG, Yi YA, Shin SJ, Park JW. Analysis of factors associated with cracked teeth. *J Endod.* 2012; 38 (3): 288-292.
- [52] Signore A, Benedicenti S, Covani U, Ravera G. A 4- to 6-year retrospective clinical study of cracked teeth restored with bonded indirect resin composite onlays. *Int J Prosthodont* .2007; 20: 609-616.

[53] Takahashi H, Finger WJ, Wegner K, Utterodt A, Komatsu M, Wöstmann B, Balkenhol M. Factors influencing marginal cavity adaptation of nanofiller containing resin composite restorations. *Dent Mater.* 2010; 26: 1166-1175.

[54] Takahashi H, Finger WJ, Endo T, Kanehira M, Koottathape N, Balkenhol M, Komatsu M. Comparative evaluation of mechanical characteristics of nanofiller containing resin composites. *American Journal of Dentistry* 2011; 24: 264-70.

[55] Watts DC, Alnazzawi A. Temperature-dependent polymerization shrinkage stress kinetics of resin-composites. *Dent Mater.* 2014; 30: 654-660.

[56] Yamamoto T, Hanabusa M, Momoi Y, Sakaguchi RL. Polymerization stress of dental resin composite continues to develop 12 hours after irradiation. *J Esthet Restor Dent.* 2015; 27: 44-54.