

**Führungsaufgaben**

- [1] Abduo J, Tennant M, McGeachie J. Lateral occlusion schemes in natural and minimally restored permanent dentition: a systematic review. *J Oral Rehabil.* 2013; 40: 788-802.
- [2] Abduo J, Tennant M. Impact of lateral occlusion schemes: A systematic review. *J Prosthet Dent.* 2015; 114: 193-204.
- [3] Al-Hiyasat AS, Abu-Alhaija ES. The relationship between static and dynamic occlusion in 14-17-year-old school children. *J Oral Rehabil.* 2004; 31: 628-633.
- [4] Attin T, Schmidlin PR, Tauböck TT. Direkte Restaurationstechnik im Seitenzahnbereich zur Versorgung von erosionsbedingtem Zahnhartsubstanzverlust. *DZZ* 2022; 77: 232-237.
- [5] Bueno MG, Tribst JPM, Borges ALS. Canine guidance reconstruction with ceramic or composite resin: A 3D finite element analysis and in vitro wear study. *J Prosthet Dent.* 2022; 127: 765.
- [6] Clark JR, Evans RD. Functional occlusion: I. A review. *J Orthod.* 2001; 28: 76-81.
- [7] Ernst CP. Die 20 beliebtesten Fehler beim Kleben. *Der Freie Zahnarzt* 2010; 11, 66-82.
- [8] Ernst, CP. Wiederauferstehung der Frasaco Stripkrone. *Dentalzeitung* 2011; 12, 14-17.
- [9] Ernst CP. Lichtpolymerisation. In: Frankenberger, R.: *Adhäsive Zahnheilkunde*, Deutscher Zahnärzte-Verlag, Köln 2013, S. 35-80.
- [10] Ernst CP. Reshaping of Incisors – An easy Solution. *Private Dentistry* 2017; 58-62.
- [11] Ernst CP. Reshaping of Incisors – An easy Solution. *Private Dentistry* 2017; 60-62.
- [12] Ernst CP. Die korrekte Vorbehandlung indirekter Restaurationen zur adhäsiven Befestigung. *ZMK* 2017; 33: 98-110.
- [13] Ernst CP, Price RB, Callaway A, Masek A, Schwarm H, Rullmann I, Willershausen B, Ehlers V. Visible Light Curing Devices - Irradiance and Use in 302 German Dental Offices. *J Adhes Dent.* 2018; 20: 41-55.
- [14] Ernst CP. Augen auf beim Lampenkauf. *ZMK* 2019; 35; 14-20.
- [15] Ernst CP. Ein praxistaugliches neues Schichtkonzept für Ceram.x Spectra ST. *ZMK* 2020; 36: 048-059.
- [16] Ernst CP. Diastemaschluss mit einer einzigen Universalfarbe. *ZMK* 2020; 36: 220-225.
- [17] Ernst CP. Komposit-Restaurationen: Differenzierte Materialauswahl bei benachbarten Kavitäten. *ZMK* 2020; 36: 599-603.
- [18] Ernst CP. Neu: etabliertes Komposit vereint universelle Farblösung mit sehr guten mechanischen Eigenschaften. *ZMK* 2020; 36: 776-780.
- [19] Ernst CP. Haltbarkeit von direkten Kompositaufbauten an Schneidezähnen nach FZT-Versorgung. *ZMK* 2021; 37: 90-102.
- [20] Ernst CP. Fallbeispiele zum ästhetischen Indikationsspektrum von Tetric Prime im Frontzahnbereich. *ZMK* 2021; 37: 542-552.

- [21] Ferracane F, Watts DC, Barghi N, Ernst CP, Rueggeberg FA, Shortall A, Price R. Der effiziente Einsatz von Lichtpolymerisationsgeräten – ein Leitfaden für Zahnärzte. ZMK 2014; 30 (4): 166-180.
- [22] FitzGerald LJ. Restoring anterior guidance by use of composite resin. Cranio. 1996; 14: 182-185.
- [23] Frankenberger R, Dudek M-C, Krämer N, Winter J, Roggendorf MJ. Die 10 beliebtesten Fehler in der Adhäsivtechnik. DZZ 2022; 77: 238-244.
- [24] Geštakovski D. The injectable composite resin technique: minimally invasive reconstruction of esthetics and function. Clinical case report with 2-year follow-up. Quintessence Int. 2019; 50:712-719.
- [25] Goldstein GR. The relationship of canine-protected occlusion to a periodontal index. J Prosthet Dent. 1979; 41: 277-283.
- [26] Hardan L, Mancino D, Bourgi R, Cuevas-Suárez CE, Lukomska-Szymanska M, Zarow M, Jakubowicz N, Zamarripa-Calderón JE, Kafa L, Etienne O, Reitzer F, Kharouf N, Haïkel Y. Treatment of Tooth Wear Using Direct or Indirect Restorations: A Systematic Review of Clinical Studies. Bioengineering 2022; 9: 346.
- [27] Heintze SD, Rousson V, Hickel R. Clinical effectiveness of direct anterior restorations - a meta-analysis. Dent Mater. 2015; 31: 481-495.
- [28] Kermanshah H, Alzwghaibi A, Al-Tufaili M, Ghabraei S. Canine Rise Method: A Conservative Approach for Worn Teeth Rehabilitation with Different Adhesive Restorative Materials. Case Rep Dent 2022; 2022: 9949879.
- [29] Kumar V, Reddy S, Kumari V S, Basha R, Mitra N. Restorative Rehabilitation of a Patient with Tooth Wear: A One-Year Clinical Follow-Up Report. Cureus. 2023; 15: e37798.
- [30] Li FF, Li M, Li M, Yang X. Modified orthodontic treatment of substitution of canines by first premolars: A case report. World J Clin Cases 2022; 10: 9078-9086.
- [31] Loomans BAC, Kreulen CM, Huijs-Visser HECE, Sterenborg BAMM, Bronkhorst EM, Huysmans MCDNJM, Opdam NJM. Clinical performance of full rehabilitations with direct composite in severe tooth wear patients: 3.5 Years results. J Dent. 2018; 70: 97-103.
- [32] Loomans B, Opdam N, Attin T et al., Severe tooth wear: European consensus statement on management guidelines. J Adhes Dent. 2017; 19: 111–119.
- [33] Lühns AK. Reparatur zahnärztlicher Seitenzahnrestaurationen – immer noch obsolet? DZZ 2015; 70: 98–109.
- [34] Mehta SB, Bronkhorst EM, Lima VP, Crins L, Bronkhorst H, Opdam NJM, Huysmans MDNJM, Loomans BAC. The effect of pre-treatment levels of tooth wear and the applied increase in the vertical dimension of occlusion (VDO) on the survival of direct resin composite restorations. J Dent. 2021; 111: 103712.
- [35] Mehta SB, Loomans BAC, van Sambeek RMF, Pereira-Cenci T, O'Toole S. Managing tooth wear with respect to quality of life: an evidence-based decision on when to intervene. Br Dent J. 2023; 234: 455-458.

- [36] Miralles R, Bull R, Manns A, Roman E. Influence of balanced occlusion and canine guidance on electromyographic activity of elevator muscles in complete denture wearers. *J Prosthet Dent.* 1989; 61: 494-498.
- [37] Murray MC, Brunton PA, Osborne-Smith K, Wilson NH. Canine risers: indications and techniques for their use. *Eur J Prosthodont Restor Dent.* 2001; 9:137-40.
- [38] Ogawa T, Ogimoto T, Koyano K. Pattern of occlusal contacts in lateral positions: canine protection and group function validity in classifying guidance patterns. *J Prosthet Dent.* 1998; 80: 67–74.
- [39] Panek H, Matthews-Brzozowska T, Nowakowska D, Panek B, Bielicki G, Makacewicz S, et al. Dynamic occlusions in natural permanent dentition. *Quintessence Int.* 2008 ;39: 337–342.
- [40] Peroz I, Leuenberg A, Haustein I, Lange KP. Comparison between balanced occlusion and canine guidance in complete denture wearers--a clinical, randomized trial. *Quintessence Int.* 2003; 34: 607-612.
- [41] Rinchuse DJ, Kandasamy S, Sciote J. A contemporary and evidence-based view of canine protected occlusion. *Am J Orthod Dentofacial Orthop.* 2007; 132: 90–102.
- [42] Shah SN, Khan S, Murtaza S, Ahmad A. Study of tooth wear in canine guidance versus group function occlusion, *J. Khyber Coll Dent.* 2014; 5: 29-32.
- [43] Sidana V, Pasricha N, Makkar M, Bhasin S. Group function occlusion. *Indian J Oral Sci.* 2012; 3: 124-128.
- [44] Thornton LJ. Anterior guidance: group function/canine guidance. A literature review. *J Prosthet Dent.* 1990; 64: 479-482.
- [45] Zhao K, Mai QQ, Wang XD, Yang W, Zhao L. Occlusal designs on masticatory ability and patient satisfaction with complete denture: a systematic review. *J Dent.* 2013; 41: 1036-1042.