

Die Guided Biofilm Therapy bleibt der absolute Favorit

[1] Blas E, Kurup AS: Introduction and methods of work. In: World Health Organization: Equity, social determinants and public health programmes. In: World Health Organization (ed). Equity, social determinants and public health programmes, 2010:3-10.

[2] World Health Organization. Political declaration of the third high-level meeting of the General Assembly on the prevention and control of noncommunicable diseases. Report by the Director-General. EXECUTIVE BOARD EB150/7, 150th session 11 January 2022, Provisional agenda item 7. 2022.

[3] Sälzer S, Graetz C, Dörfer CE, et al.: Contemporary practices for mechanical oral hygiene to prevent periodontal disease. *Periodontology 2000*. 2020;84(1):35-44.

<https://pubmed.ncbi.nlm.nih.gov/32844413/>

<https://onlinelibrary.wiley.com/doi/pdfdirect/10.1111/prd.12332>

[4] Axelsson P, Nystrom B, Lindhe J.: The long-term effect of a plaque control program on tooth mortality, caries and periodontal disease in adults. Results after 30 years of maintenance. *J Clin Periodontol*. 2004;31(9):749-57. Epub 2004/08/18. <https://pubmed.ncbi.nlm.nih.gov/15312097/>

[5] Tonetti MS, Chapple ILC, Jepsen S, et al.: Primary and secondary prevention of periodontal and peri-implant diseases. *Journal of Clinical Periodontology*. 2015;42:S1-S4.

<https://pubmed.ncbi.nlm.nih.gov/25683242/>

[6] Bastendorf K, Strafela-Bastendorf N.: Auf das klinische Protokoll kommt es an – PZR, UPT und GBT. *Quintessenz*. 2020;71(12):1380-9.

[7] Vouros I, Antonoglou GN, Anoixiadou S, et al.: A novel biofilm removal approach (Guided Biofilm Therapy) utilizing erythritol air-polishing and ultrasonic piezo instrumentation: A randomized controlled trial. *Int J Dent Hyg*. 2021;n/a(n/a). Epub 2021/07/05.

<https://www.ncbi.nlm.nih.gov/pubmed/34218516>

[8] Bischoff JG.: Herausforderung Prophylaxezentrum. *Zahnarzt Wirtschaft Praxis*. 2018(6):16-9.

[9] Bühler J, Amato M, Weiger R, et al.: A systematic review on the patient perception of periodontal treatment using air polishing devices. *Int J Dent Hyg*. 2016;14(1):4-14. Epub 2015/01/27.

<https://www.ncbi.nlm.nih.gov/pubmed/25619863>

[10] Bühler J, Amato M, Weiger R, et al.: A systematic review on the effects of air polishing devices on oral tissues. *Int J Dent Hyg*. 2016;14(1):15-28. Epub 2015/02/19.

<https://www.ncbi.nlm.nih.gov/pubmed/25690301>

<https://onlinelibrary.wiley.com/doi/10.1111/idh.12120>

[11] Fu JH, Wong LB, Tong HJ, et al.: Conventional versus comprehensive dental prophylaxis: comparing the clinical outcomes between rubber cup and air polishing and the importance of plaque disclosure. *Quintessence Int*. 2021;0(0):0. Epub 2021/01/26.

<https://pubmed.ncbi.nlm.nih.gov/33491396/>

[12] Lang N, A. L, KD. B.: Wissenschaftlicher Konsensus Guided Biofilm Therapy-Protokoll. Ein neues Konzept für die primäre und sekundäre Prävention. 2019. [\[English version available\]](#)

[13] Arefnia B, Koller M, Wimmer G, et al.: In Vitro Study of Surface Changes Induced on Enamel and Cementum by Different Scaling and Polishing Techniques. Oral Health Prev Dent. 2021;19(1):85-92. Epub 2021/01/30. <https://www.ncbi.nlm.nih.gov/pubmed/33511822>

[14] Burkhardt AS, et al.: Effect of air-polishing using erythritol on surface roughness of enamel and dentine compared to conventional methods. Poster presented at the EuroPerio, Copenhagen 2022. 2022.

[15] Stiftung Warentest. Weit aufmachen, bitte. Stiftung Warentest: Professionelle Zahnreinigung. test. 2015(7):86-90.

[16] Mensi M, Scotti E, Sordillo A, et al.: Plaque disclosing agent as a guide for professional biofilm removal: A randomized controlled clinical trial. Int J Dent Hyg. 2020;18(3):285-94. Epub 2020/04/30. <https://www.ncbi.nlm.nih.gov/pubmed/32348624>

[17] Donnet M, Fournier M, Schmidlin PR, et al. : A Novel Method to Measure the Powder Consumption of Dental Air-Polishing Devices. Applied Sciences. 2021;11(3):1101.

<https://www.mdpi.com/2076-3417/11/3/1101>

[18] Koch JH. Die „Guided Biofilm Therapy“: ist der absolute Favorit bei den Patienten. Weltweite Umfrage zu bevorzugter Prophylaxe-Methode. ZMK. 2022;38(4):183-5. [\[English version available\]](#)

[19] Feil PH, Grauer JS, Gadbury-Amyot CC, et al.: Intentional use of the Hawthorne effect to improve oral hygiene compliance in orthodontic patients. J Dent Educ. 2002;66(10):1129-35. <https://www.ncbi.nlm.nih.gov/pubmed/12449206>

[20] Furrer C, Battig R, Votta I, et al.: Patientenakzeptanz nach Umstellung auf «Guided Biofilm Therapy». Swiss Dent J. 2021;131(3):229-34. <https://pubmed.ncbi.nlm.nih.gov/33666383/> [\[English version available\]](#)