

- [1] Augusto MG, Torres C, Pucci CR, Schlueter N, Borges AB: Bond stability of an universal adhesive system to eroded/abraded dentin after deproteinization. *Oper Dent* 43, 291–300 (2018).
- [2] Azzopardi A, Bartlett DW, Watson TF, Sherriff M: The measurement and prevention of erosion and abrasion. *J Dent* 29, 395–400 (2001).
- [3] Bartlett D, Sundaram G, Moazzez R: Trial of protective effect of fissure sealants, in vivo, on the palatal surfaces of anterior teeth, in patients suffering from erosion. *J Dent* 39, 26–29 (2011).
- [4] Carvalho TS, Colon P, Ganss C, Huysmans MC, Lussi A, Schlueter N et al.: Consensus Report of the European Federation of Conservative Dentistry: Erosive tooth wear diagnosis and management. *Swiss Dent J* 126, 342–346 (2016).
- [5] Deari S, Wegehaupt FJ, Taubock TT, Attin T: Influence of different pretreatments on the microtensile bond strength to eroded dentin. *J Adhes Dent* 19, 147–155 (2017).
- [6] Flury S, Koch T, Peutzfeldt A, Lussi A, Ganss C: The effect of a tin-containing fluoride mouth rinse on the bond between resin composite and erosively demineralised dentin. *Clin Oral Investig* 17, 217–225 (2013).
- [7] Ganss C, Hardt M, Lussi A, Cocks AK, Klimek J, Schlueter N: Mechanism of action of tin-containing fluoride solutions as anti-erosive agents in dentine – an in vitro tin-uptake, tissue loss, and scanning electron microscopy study. *Eur J Oral Sci* 118, 376–384 (2010).
- [8] Ganss C, Lussi A, Peutzfeldt A, Naguib Attia N, Schlueter N: Impact of Sn/F pre-treatments on the durability of protective coatings against dentine erosion/abrasion. *PLoS One* 10, e0123889 (2015).
- [9] Ganss C, Neutard L, von Hinckeldey J, Klimek J, Schlueter N: Efficacy of a tin/fluoride rinse: a randomized in situ trial on erosion. *J Dent Res* 89, 1214–1218 (2010).
- [10] Hirce JD, Sather AH, Chao EY: The effect of topical fluorides, after acid etching of enamel, on the bond strength of directly bonded orthodontic brackets. *Am J Orthod* 78, 444–452 (1980).
- [11] Huysmans MC, Young A, Ganss C: The role of fluoride in erosion therapy. *Monogr Oral Sci* 25, 230–243 (2014).
- [12] Imamura N, Kawaguchi T, Shimizu H, Takahashi Y: Effect of three metal priming agents on the bond strength of adhesive resin cement to Ag-Zn-Sn-In alloy and component metals. *Dent Mater J* 37, 301–307 (2018).
- [13] Loomans B, Opdam N, Attin T, Bartlett D, Edelhoff D, Frankenberger R et al.: Severe tooth wear: European Consensus Statement on Management Guidelines. *J Adhes Dent* 19, 111–119 (2017).
- [14] Low T, Von Fraunhofer JA, Winter GB: The bonding of a polymeric fissure sealant to topical fluoride-treated teeth. *J Oral Rehabil* 2, 303–307 (1975).
- [15] Low T, von Fraunhofer JA, Winter GB: Influence of the topical application of fluoride on the in vitro adhesion of fissure sealants. *J Dent Res* 56, 17–20 (1977).
- [16] Oliveira B, Ulbaldini A, Sato F, Baesso ML, Bento AC, Andrade L et al.: Chemical interaction analysis of an adhesivecontaining 10-methacryloyloxydecyl dihydrogen phosphate (10-MDP) with the dentin in noncarious cervical lesions. *Oper Dent* 42, 357–366 (2017).
- [17] Petzold M: The influence of different fluoride compounds and treatment conditions on

- dental enamel: a descriptive in vitro study of the CaF₂ precipitation and microstructure. *Caries Res* 35 (Suppl 1), 45–51 (2001).
- [18] Peumans M, De Munck J, Van Landuyt K, Van Meerbeek B: Thirteen-year randomized controlled clinical trial of a two-step self-etch adhesive in non-cariou cervical lesions. *Dent Mater* 31, 308–314 (2015).
- [19] Peutzfeldt A, Jaeggi T, Lussi A: Restorative therapy of erosive lesions. *Monogr Oral Sci* 25, 253–261 (2014).
- [20] Peutzfeldt A, Koch T, Ganss C, Flury S, Lussi A: Effect of tin-chloride pretreatment on bond strength of two adhesive systems to dentin. *Clin Oral Investig* 18, 535–543 (2014).
- [21] Saxegaard E, Rolla G: Fluoride acquisition on and in human enamel during topical application in vitro. *Scand J Dent Res* 96, 523–535 (1988).
- [22] Schlueter N, Hardt M, Lussi A, Engelmann F, Klimek J, Ganss C: Tin-containing fluoride solutions as anti-erosive agents in enamel: an in vitro tin-uptake, tissue-loss, and scanning electron micrograph study. *Eur J Oral Sci* 117, 427–434 (2009).
- [23] Schlueter N, Peutzfeldt A, Ganss C, Lussi A: Does tin pre-treatment enhance the bond strength of adhesive systems to enamel? *J Dent* 41, 642–652 (2013).
- [24] Schlüter N, Gressbach C, Ziemann C, Winterfeld T, Ganß C: Kompositbasierte Beschichtungen zur Prävention von Dentinerosionen. *Dtsch Zahnärztl Z* 69, 564–572 (2014).
- [25] Siqueira F, Cardenas A, Gomes GM, Chibinski AC, Gomes O, Bandeca MC et al.: Three-year effects of deproteinization on the in vitro durability of resin/dentin-eroded interfaces. *Oper Dent* 43, 60–70 (2018).
- [26] Umwali A, Askar H, Paris S, Schwendicke F: Radiographic, antibacterial and bond-strength effects of radiopaque caries tagging. *Sci Rep* 6, 27319 (2016).
- [27] Yetkiner E, Wegehaupt FJ, Attin R, Wiegand A, Attin T: Stability of two resin combinations used as sealants against toothbrush abrasion and acid challenge in vitro. *Acta Odontol Scand* 72, 825–830 (2014).
- [28] Zimmerli B, De Munck J, Lussi A, Lambrechts P, Van Meerbeek B: Long-term bonding to eroded dentin requires superficial bur preparation. *Clin Oral Investig* 16, 1451–1461 (2012).