

ZMK 34(10) 2018, S. 656-662

Prof. Dr. med. dent. habil. Christian R. Gernhardt, Henrike Jäger

Endodontie 2018: Was geht, was bleibt, was kommt?

- [1] Friedman S, Mor C: The success of endodontic therapy-healing and functionality. J Calif Dent Assoc 32 (6), 493–503 (2004).
- [2] Ng YL, Mann V, Gulabivala K: Tooth survival following non-surgical root canal treatment: a systematic review of the literature. Int Endod J 43 (3), 171–189 (2010).
- [3] Azim AA, Griggs JA, Huang GT: The Tennessee Study: factors affecting treatment outcome and healing time following nonsurgical root canal treatment. Int Endod J 49, 6–16 (2016).
- [4] Gillen BM et al.: Impact of the quality of coronal restoration versus the quality of root canal fillings on success of root canal treatment: a systematic review and meta-analysis. J Endod 37 (7), 895–902 (2011).
- [5] Siqueira jr. J: Aetiology of root canal failure: why well-treated teeth can fail. Int Endod J 34, 1–10 (2001).
- [6] Weber M, Stratz N, Hannig C: Möglichkeiten und Grenzen der Darstellung von endodontischen Strukturen mit DVT: Swiss Dental Journal SSO 125, 303–311 (2015).
- [7] European Society of Endodontology, 2006.
- [8] Walia HM, Brantley WA, Gerstein H: An initial investigation of the bending and torsional properties of Nitinol root canal files. J Endodont 14, 346–351 (1988). [Thompson, 2000.](#)
- [9] Pereira WS et al.: Metallurgical characterization of M-Wire nickel-titanium shape memory alloy used for endodontic rotary instruments during low-cycle fatigue. Journal of Endodontics 38 (1), 105–107 (2012).
- [10] Bürklein S, Benten S, Schäfer E: Shaping ability of different single-file systems in severely curved root canals of extracted teeth. IEJ 46 (6), 590–597 (2012).
- [11] Schloss T, Wollner J: Modernste Endodontie mit einem einfachen Konzept. Endo Tribune German Edition 11 (2013).
- [12] Pedula E, Grande NM, Plotino G, Gambarini G, Rapisarda E: Influence of continuous or reciprocating motion on cyclic fatigue resistance of 4 different nickel-titanium rotary instruments. Journal of Endodontics 39 (2), 258–261 (2013).
- [13] Metzger Z et al.: The quality of root canal preparation and root canal obturation in canals treated with rotary *versus* self-adjusting files: a three-dimensional micro-computed tomographic study. Journal of Endodontics 36 (9), 1569–1573 (2010).
- [14] Peters OA, Schönenberger K, Laib A: Effects of four Ni-Ti preparation techniques on root canal geometry assessed by micro computed tomography. Int Endod J 34 (3), 221–230 (2001).
- [15] Hülsmann M: Checklisten der Zahnmedizin: Endodontie, Desinfektion des endodontischen Systems. Stuttgart, New York, Delhi, Rio: Thieme, 147 (2008).
- [16] Gopikrishna V, Pare S, Pradeep Kumar A, Lakshmi Narayanan L: Irrigation protocol among endodontic faculty and post-graduate students in dental colleges of India: a survey. J Conserv Dent 16, 394–398 (2013).
- [17] Zehnder M: Root canal irrigants. J Endod 32, 389–398 (2006).
- [18] Camara AC, de Albuquerque MM, Aguiar CM, de Barros Correia AC: In vitro antimicrobial activity of 0,5%, 1%, and 2,5% sodium hypochlorite in root canals instrumented with the ProTaper Universal system. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 108, 55–61 (2009).

Kommentar [AK1]: ?? Weitere Literaturstelle?

- [19] Jose J, Krishnamma S, Peedikayil F, Aman S, Tomy N, Mariodan JP: Comparative evaluation of antimicrobial activity of QMiX, 2,5% sodium hypochlorite, 2% chlorhexidine, guava leaf extract and aloe vera extract against *Enterococcus faecalis* and *Candida albicans* – an in-vitro study. J Clin Diagn Res 10, ZC20–23 (2016).
- [20] Cullen JK, Wealleans JA, Kirkpatrick TC, Yaccino JM: The effect of 8,25% sodium hypochlorite on dental pulp dissolution and dentin flexural strength and modulus. J Endod 41, 920–924 (2015).
- [21] Dumitriu D, Dobre T: Effects of temperature and hypochlorite concentration on the rate of collagen dissolution. J Endod 41, 903–906 (2015).
- [22] Vaghela DJ, Kandaswamy D, Venkateshbabu N, Jamini N, Ganesh A: Disinfection of dentinal tubules with two different formulations of calcium hydroxide as compared to 2 % chlorhexidine: as intracanal medicaments against *Enterococcus faecalis* and *Candida albicans*: an in vitro study. J Conserv Dent 14, 182–186 (2011).
- [23] Mahendra A, Koul M, Upadhyay V, Dwivedi R: Comparative evaluation of antimicrobial substantivity of different concentrations of chlorhexidine as a root canal irrigant: an in vitro study. J Oral Biol Craniofac Res 4, 181–185 (2014).
- [24] Carrilho MR, Carvalho RM, Sousa EN et al.: Substantivity of chlorhexidine to human dentin. Dent Mater 26, 779–785 (2010).
- [25] Kandaswamy D, Venkateshbabu N, Gogulnath D, Kindo AJ: Dentinal tubule disinfection with 2% chlorhexidine gel, propolis, morinda citrifolia juice, 2% povidone iodine, and calcium hydroxide. Int Endod J 43, 419–423 (2010).
- [26] Schmidt TF, Teixeira CS, Felipe MC, Felipe WT, Pashley DH, Bortoluzzi EA: Effect of ultrasonic activation of irrigants on smear layer removal. J Endod 41, 1359–1363 (2015).
- [27] Urban K, Donnermeyer D, Schafer E, Burklein S: Canal cleanliness using different irrigation activation systems: a SEM evaluation. Clin Oral Investig 21, 2681–2687 (2017).
- [28] AAE Clinical Considerations for a Regenerative Procedure Revised 4/1/2018.
- [29] Taneja S, Kumari M: Use of triple antibiotic paste in the treatment of large periradicular lesions. Journal of Investigative and Clinical Dentistry 3 (1), 72–76 (2012).
- [30] Kim J-H, Kim Y, Shin S-J, Park J-W, Jung I-Y: Tooth discoloration of immature permanent incisor associated with triple antibiotic therapy: a case report. Journal of Endodontics 36 (6), 1086–1091 (2010).