

Vorhersagbarkeit in der Endodontie – Diagnostik und Grundlagen
Dr. Andreas Habash
ZMK 11 (33), S. 814-819

- [1] Krasner P1, Rankow HJ.: Anatomy of the pulp-chamber floor. J Endod. 2004 Jan;30(1):5-16.
- [2] Juloski J1, Radovic I, Goracci C, Vulicevic ZR, Ferrari M. Ferrule effect: a literature review. J Endod. 2012 Jan;38(1):11-9.
- [3] Arnold M, Friedrichs C, Tulus G, Verch S, Dennhardt H, Sanner F. Intrakoronale und intrakanal.re endodontische Diagnostik (IKD). Endodontie 2013; 22:9-21.
- [4] . Siqueira JF Jr1, Rôças IN, Favieri A, Lima KC.: Chemomechanical reduction of the bacterial population in the root canal after instrumentation and irrigation with 1%, 2.5%, and 5.25% sodium hypochlorite J Endod. 2000 Jun;26(6):331-4.
- [5] Gurgel-Filho ED1, Vivacqua-Gomes N, Gomes BP, Ferraz CC, Zaia AA, Souza-Filho FJ.: In vitro evaluation of the effectiveness of the chemomechanical preparation against *Enterococcus faecalis* after single- or multiple-visit root canal treatment. Braz Oral Res. 2007 Oct-Dec;21(4):308-13.:
- [6] van der Sluis L., Boutsoukis C., Jiang LM., Macedo R., Verhaagen B., Versluis M. (2015) Root Canal Irrigation. In: Chávez de Paz L., Sedgley C., Kishen A. (eds) The Root Canal Biofilm. Springer Series on Biofilms, vol 9. Springer, Berlin, Heidelberg
- [7] David E. Jaramillo, Enrique Aguilar, Ana Arias, Ronald Ordinola-Zapata, Raydolfo M. Aprecio and Jose L. Ibarrola: Root canal disinfection comparing conventional irrigation vs photon-induced photoacoustic streaming (PIPS) using a buffered 0.5 % sodium hypochlorite solution. The purpose of this study was to assess the antimicrobial effect of a buffered 0.5 % sodium hypochlorite solution activated by photon-induced photoacoustic streaming compared to conventional irrigation