

ZMK 6/2022 (38), 284-287

Dr. Paul Leonhard Schuh, PD Dr. Marcel Hanisch

Weichgewebsmanagement in der ästhetischen Zone

[1] Cardaropoli G, Araújo M, Lindhe J. Dynamics of bone tissue formation in tooth extraction sites. An experimental study in dogs. *J Clin Periodontol.* 2003; 30 (9): 809-18.

[2] Wang J, Wang L, Zhou Z, Lai H, Xu P, Liao L, Wei J. Biodegradable Polymer Membranes Applied in Guided Bone/Tissue Regeneration: A Review. *Polymers* 2016; 8, 115.

[3] Atieh MA, Alsabeeha NHM. Soft tissue changes after connective tissue grafts around immediately placed and restored dental implants in the esthetic zone: A systematic review and meta-analysis. *J Esthet Restor Dent.* 2020; 32 (3): 280-290.

[4] Cairo F, Barbato L, Selvaggi F, Baielli MG, Piattelli A, Chambrone L. Surgical procedures for soft tissue augmentation at implant sites. A systematic review and meta-analysis of randomized controlled trials. *Clin Implant Dent Relat Res.* 2019; 21 (6): 1262-1270.

[5] Thoma DS, Buranawat B, Hammerle CH, Held U, Jung RE. Efficacy of soft tissue augmentation around dental implants and in partially edentulous areas: A systematic review. *Journal of Clinical Periodontology*, 2014; 41 (Suppl 15), S77–S91.

[6] Puisys A, Linkevicius T. The influence of mucosal tissue thickening on crestal bone stability around bone-level implants. A prospective controlled clinical trial. *Clin Oral Implants Res.* 2015; 26 (2): 123-9.

[7] Schuh PL, Wachte H, Bolz W, Maischberger C, Schenk A, Kühn M. „Teflon-Tape-Technique“: synergy between isolation and lucidity. *Quintessenz International* Volume 2019; 50 (1).