

- [1] J. J. Hyman and W. Doblecki, "Computerized endodontic diagnosis," *J. Am. Dent. Assoc.* 1939, vol. 107, no. 5, pp. 755–758, Nov. 1983, doi: 10.14219/jada.archive.1983.0349.
- [2] J. E. Overall and J. M. Williams, "Conditional probability program for diagnosis of thyroid function," *JAMA*, vol. 183, pp. 307–313, Feb. 1963, doi: 10.1001/jama.1963.03700050039009.
- [3] S. Weiss, C. A. Kulikowski, and A. Safir, "Glaucoma consultation by computer," *Comput. Biol. Med.*, vol. 8, no. 1, pp. 25–40, Jan. 1978, doi: 10.1016/0010-4825(78)90011-2.
- [4] L.-L. Kirkevang, D. Ørstavik, G. Bahrami, A. Wenzel, and M. Vaeth, "Prediction of periapical status and tooth extraction," *Int. Endod. J.*, vol. 50, no. 1, pp. 5–14, Jan. 2017, doi: 10.1111/iej.12581.
- [5] S. Sadr *et al.*, "Deep Learning for Detection of Periapical Radiolucent Lesions: A Systematic Review and Meta-analysis of Diagnostic Test Accuracy.," *J. Endod.*, pp. S0099-2399(22)00845–7, Dec. 2022, doi: 10.1016/j.joen.2022.12.007.
- [6] S. R. Herbst, V. Pitchika, J. Krois, A. Krasowski, and F. Schwendicke, "Machine Learning to Predict Apical Lesions: A Cross-Sectional and Model Development Study," *J. Clin. Med.*, vol. 12, no. 17, p. 5464, Aug. 2023, doi: 10.3390/jcm12175464.
- [7] S. Patel, E. Brady, R. Wilson, J. Brown, and F. Mannocci, "The detection of vertical root fractures in root filled teeth with periapical radiographs and CBCT scans," *Int. Endod. J.*, vol. 46, no. 12, pp. 1140–1152, Dec. 2013, doi: 10.1111/iej.12109.
- [8] M. Edlund, M. K. Nair, and U. P. Nair, "Detection of vertical root fractures by using cone-beam computed tomography: a clinical study," *J. Endod.*, vol. 37, no. 6, pp. 768–772, Jun. 2011, doi: 10.1016/j.joen.2011.02.034.
- [9] S. Kositbowornchai, S. Plermkamon, and T. Tangkosol, "Performance of an artificial neural network for vertical root fracture detection: an *ex vivo* study," *Dent. Traumatol.*, vol. 29, no. 2, pp. 151–155, Apr. 2013, doi: 10.1111/j.1600-9657.2012.01148.x.
- [10] M. Johari, F. Esmaeili, A. Andalib, S. Garjani, and H. Saberhari, "Detection of vertical root fractures in intact and endodontically treated premolar teeth by designing a probabilistic neural network: an *ex vivo* study," *Dentomaxillofacial Radiol.*, vol. 46, no. 2, p. 20160107, Feb. 2017, doi: 10.1259/dmfr.20160107.
- [11] X. Zhang *et al.*, "A Cone-Beam Computed Tomographic Study on Mandibular First Molars in a Chinese Subpopulation," *PLOS ONE*, vol. 10, no. 8, p. e0134919, Aug. 2015, doi: 10.1371/journal.pone.0134919.
- [12] S.-J. Jeon *et al.*, "Deep-learning for predicting C-shaped canals in mandibular second molars on panoramic radiographs," *Dento Maxillo Facial Radiol.*, vol. 50, no. 5, p. 20200513, Jul. 2021, doi: 10.1259/dmfr.20200513.
- [13] S. Yang *et al.*, "Development and Validation of a Visually Explainable Deep Learning Model for Classification of C-shaped Canals of the Mandibular Second Molars in Periapical and Panoramic Dental Radiographs," *J. Endod.*, vol. 48, no. 7, pp. 914–921, Jul. 2022, doi: 10.1016/j.joen.2022.04.007.
- [14] C. Buyuk, B. Arican Alpay, and F. Er, "Detection of the separated root canal instrument on panoramic radiograph: a comparison of LSTM and CNN deep learning methods.," *Dento Maxillo Facial Radiol.*, p. 20220209, Jan. 2023, doi: 10.1259/dmfr.20220209.
- [15] O. Amza, B. Dimitriu, I. Suci, R. Bartok, and M. Chirila, "Etiology and Prevention of an Endodontic Iatrogenic Event: Instrument Fracture," *J. Med. Life*, vol. 13, no. 3, pp. 378–381, 2020, doi: 10.25122/jml-2020-0137.
- [16] L. Albitar, T. Zhao, C. Huang, and M. Mahdian, "Artificial Intelligence (AI) for Detection and Localization of Unobturated Second Mesial Buccal (MB2) Canals in Cone-Beam Computed Tomography (CBCT).," *Diagn. Basel Switz.*, vol. 12, no. 12, Dec. 2022, doi: 10.3390/diagnostics12123214.
- [17] American Association of Endodontists AAE, "AAE Endodontic Case Difficulty Assessment Form

and Guidelines." 2022.

[18] S. R. Haug, A. F. Solfjeld, L. E. Ranheim, and A. Bårdsen, "Impact of Case Difficulty on Endodontic Mishaps in an Undergraduate Student Clinic," *J. Endod.*, vol. 44, no. 7, pp. 1088–1095, 2018, doi: 10.1016/j.joen.2018.03.012.

[19] S. R. Herbst, C. S. Herbst, and F. Schwendicke, "Preoperative risk assessment does not allow to predict root filling length using machine learning: A longitudinal study," *J. Dent.*, vol. 128, p. 104378, Jan. 2023, doi: 10.1016/j.jdent.2022.104378.

[20] C. S. Herbst, F. Schwendicke, J. Krois, and S. R. Herbst, "Association between patient-, tooth- and treatment-level factors and root canal treatment failure: A retrospective longitudinal and machine learning study," *J. Dent.*, vol. 117, p. 103937, Feb. 2022, doi: 10.1016/j.jdent.2021.103937.

[21] J. Lee *et al.*, "An Endodontic forecasting model based on the analysis of preoperative dental radiographs: A pilot study on an endodontic predictive deep neural network," *J. Endod.*, pp. S0099-2399(23)00178–4, Apr. 2023, doi: 10.1016/j.joen.2023.03.015.

[22] D. Helvacioğlu-Yigit and A. Sinanoğlu, "Use of cone-beam computed tomography to evaluate C-shaped root canal systems in mandibular second molars in a Turkish subpopulation: a retrospective study," *Int. Endod. J.*, vol. 46, no. 11, pp. 1032–1038, Nov. 2013, doi: 10.1111/iej.12094.

[23] F. Schwendicke, S. Mertens, A. G. Cantu, A. Chaurasia, H. Meyer-Lueckel, and J. Krois, "Cost-effectiveness of AI for caries detection: randomized trial," *J. Dent.*, vol. 119, p. 104080, Apr. 2022, doi: 10.1016/j.jdent.2022.104080.

Abbildung 1 Begriffsbestimmung der künstlichen Intelligenz

Abbildung 2 Durch