

Nezar Watted, Muhamad Abu-Hussein, Obaida Awadi, Peter Proff, Borbély Péter, Viktória Hegedűs

Ist die approximale Schmelzreduktion – Stripping – eine Behandlungsmethode für Platzbeschaffung in der Kieferorthopädie?

- [1] Alexandra RG: The Alexandra discipline contemporary concepts and philosophies. Gray A. *Angel Ormco Corporation*, 1986, 461 pages, 353 illustrations
- [2] Ballard ML: Asymmetry in tooth size: a factor in the etiology – diagnosis and treatment of mal-occlusion. *Angle Orthod* 14, 67–70 (1944).
- [3] Barrer HG: Protecting the integrity of mandibular incisor position through keystoneing procedure and spring retainer appliance. *J Clin Orthod* 9, 486–494 (1975).
- [4] Betteridge MA: The effects of interdental strip ping on labial segments evaluated one year out of retention. *Br J Orthod* 8, 193–197 (1981).
- [5] Begg PR: Stone age man's dentition. *Am J Orthod* 4, 298–531 (1954).
- [6] Boese LR: Fiberotomy and reproximation without lower retention, nine years in retrospect. *Angel Orthod Part I* 50, 88–97 (1980), part II 50,169–178 (1980).
- [7] Bolton WA: Disharmony in tooth size and its relation to the analysis and treatment of malocclusion. *Angle Orthod* 28,113–130 (1958).
- [8] Carey CW: Linear arch dimension and tooth size. *Am J Orthod* 35, 762–769 (1949).
- [9] Crain G, Sheriden JJ: Susceptibility to caries and periodontal disease after posterior air-rotor stripping, *J Clin Orthod* 24, 84–85 (1990).
- [10] Fillion D: Zur approximalen Schmelzreduktion in der Erwachsenenkieferorthopädie. Teil I: Anatomische und parodontale Gesichtspunkte. *Inf Orthod Kieferthop* 27, 48–63 (1995).
- [11] Fillion D: Zur approximalen Schmelzreduktion in der Erwachsenenkieferorthopädie. Teil II: Vor- und Nachteile der approximalen Schmelzreduktion. *Inf Orthod Kieferthop* 27, 64–90 (1995).
- [12] Fillion D: Zur approximalen Schmelzreduktion in der Erwachsenenkieferorthopädie. Teil III: Grundprinzipien der approximalen Schmelzreduktion und Vorgehensweise anhand dreier Fälle. *Inf Orthod Kieferthop* 27, 91–105 (1995).
- [13] Gillings BM, Buonocore M: An investigation of enamel thickness in human lower incisor teeth. *J Dent Res* 40,105–118 (1961).
- [14] Giovannoli JL, Heins PJ: Classification de l'aspect radiographique des lésions osseuses en rapport avec une proximité radiculaire. *J Parodontol* 2, 31–40 (1983).
- [15] Heins PJ, Wieder SM: A histological study of the width and nature of interradicular space in human adult premolars and molars. *J Dent Res* 65, 948–951 (1986).
- [16] Hudson AL: A study of the effects of mesio-distal reduction of mandibular anterior teeth. *Am J Orthod* 42, 615–624 (1956).

- [17] Joseph VP, Rossouw PE, Basson NJ: Orthodontic microabrasive reoximation. *Am J Orthod Dentofacial Orthod* 102, 351–359 (1992).
- [18] Mangourey NH, Moussa MM, Mostafa YA, Girgis AS: In vivo remineralization after air-rotor stripping. *J Clin Orthod* 25, 75–78 (1991).
- [19] Paskow H: Self-alignment following interproximal stripping. *Am J Ortod* 58, 240–249 (1970).
- [20] Peck H, Peck S: An index for assessing tooth shape deviations as applied to the mandibular incisors. *Am J Orthod* 61, 384–401 (1972).
- [21] Peck S, Peck L, Kataja M: The gingival smile line. *Angle Orthod* 62, 91–100 (1992).
- [22] Sadowsky C, Begole EA: Long term effects of orthodontic treatment on periodontal health. *Am J Orthod* 80, 156–172 (1981).
- [23] Sheridan JJ: Air-rotor stripping. *J Clin Orthod* 19, 43–59 (1985).
- [24] Sheridan JJ: Air-rotor stripping update. *J Clin Orthod* 21, 781–788 (1987).
- [25] Shillingburg HT, Grace CS: Thickness of enamel and dentin. *JS Calif St Dent Assoc* 4, 33–52 (1973).
- [26] Takei HH: The interdental space. *Dent Clin North Am* 24, 169–176 (1980).
- [27] Tarnow D, Magner A, Fletcher P: The effect of the distance from the contact point to the crest of bone on the presence or absence of the interproximal dental papilla. *J Periodontol* 63, 995–996 (1992).
- [28] Tallec P: Anatomie dentaire (352 p.). I.F.O.S.U.P.D. éd., (1986). Gedrucktes Buch.
- [29] Tuverson DL: Anterior interocclusal relations: part I. *Am J Orthod* 78, 361–370 (1980).
- [30] Watted N, Teuscher T, Wieber M: Die interdisziplinäre Zusammenarbeit in der Zahnmedizin: Präprothetische Kieferorthopädie. *Quintessenz* 52 (5), 463–476 (2001).
- [31] Watted N, Teuscher T: Die interdisziplinäre Zusammenarbeit zwischen Kieferorthopädie und Parodontologie. *Kieferorthop* 16, 103–119 (2002).
- [32] Watted N, Teuscher T: interdisziplinäre Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik, Parodontologie, Orthodontie und Prothetik. *ZMK* 18 (5), 304–315 (2002).
- [33] Watted N, Teuscher T, Ries St: Die interdisziplinäre Behandlung und dentofazialen Ästhetik. *ZMK* 19 (11), 744–754 (2003).
- [34] Watted N, Abdulgani Azz, Abu-Hussein M: Aesthetic replacement of congenitally missing tooth using Fober-Reinforced Composite (FRC). *Journal of Dental and Health Sciences (IJDHS)* 4 (1), 430–439 (2014).
- [35] Watted N, Borbely P, Watted A, Ghannam N, Abdulgani E, Abu-Hussein M: Multidisciplinary approach in the rehabilitation of congenitally missing lateral incisors: a new trend in daily practice. *Oral Health and Care* 1 (1), 2–8 (2016).
- [36] Watted N, Borbely P, Watted A, Ghannam N, Abudalgani E, Abu-Hussein M: Multidisciplinary approach in the rehabilitation of congenitally missing lateral incisors: a new trend in daily practice. *Oral Health and Care* 1 (1), 1–8 (2017).

[37] Wheeler RC: Dental anatomy, physiology and occlusion (520 p.). W.B. Saunders Co, Philadelphia (1974).

[38] Williams R: Eliminating lower retention. J Clin Orthod 19, 342–348 (1985).

[39] Zachrisson BU, Alnaes L: Periodontal condition in orthodontically treated and untreated individuals. I: Loss of attachment, gingival pocket depth and clinical crown height. Angle Orthod 43, 402–411 (1973). II: Alveolar bone loss: Radiographic findings. 44, 48–55 (1974).