

Nichtchirurgische Revisionsbehandlung

Dr. Günther Stöckl

ZMK 33 (11) 2017, S. 820-824

- [1] Alley BS, et al: A comparison of survival of teeth following endodontic treatment performed by general dentists or by specialists. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 98 (1), 115-8 (2004).
- [2] Iqbal MK, Kim S: For teeth requiring endodontic treatment, what are the differences in outcomes of restored endodontically treated teeth compared to implant-supported restorations? *Int J Oral Maxillofac Implants* 22 Suppl, 96-116 (2007).
- [3] Doyle SL, et al: Retrospective cross sectional comparison of initial nonsurgical endodontic treatment and single-tooth implants. *J Endod* 32 (9), 822-7 (2006).
- [4] Kirkevang LL, et al: Frequency and distribution of endodontically treated teeth and apical periodontitis in an urban Danish population. *Int Endod J* 34 (3), 198-205 (2001).
- [5] Kirkevang LL, et al: Risk factors for developing apical periodontitis in a general population. *Int Endod J* 40 (4), 290-9 (2007).
- [6] Ng YL, Mann V, Gulabivala K: A prospective study of the factors affecting outcomes of nonsurgical root canal treatment: part 1: periapical health. *Int Endod J* 44 (7), 583-609 (2011).
- [7] Torabinejad M, et al: Outcomes of nonsurgical retreatment and endodontic surgery: a systematic review. *J Endod* 35 (7), 930-7 (2009).
- [8] He J, et al: Clinical and Patient-centered Outcomes of Nonsurgical Root Canal Retreatment in First Molars Using Contemporary Techniques. *J Endod* 43 (2), 231-237 (2017).
- [9] Ng YL, Mann V, Gulabivala K: Outcome of secondary root canal treatment: a systematic review of the literature. *Int Endod J* 41 (12), 1026-46 (2008).
- [10] de Chevigny C, et al: Treatment outcome in endodontics: the Toronto study--phases 3 and 4: orthograde retreatment. *J Endod* 34 (2), 131-7 (2008).
- [11] Gorduysus MO, Gorduysus M, Friedman S: Operating microscope improves negotiation of second mesiobuccal canals in maxillary molars. *J Endod* 27 (11), 683-6 (2001).
- [12] Arias MP, et al: Effect of ultrasonic streaming on intra-dental disinfection and penetration of calcium hydroxide paste in endodontic treatment. *J Appl Oral Sci* 24 (6), 575-581 (2016).
- [13] Shahabinejad H, et al: Success of ultrasonic technique in removing fractured rotary nickel-titanium endodontic instruments from root canals and its effect on the required force for root fracture. *J Endod* 39 (6), 824-8 (2013).
- [14] Conde AJ, et al: Effect of sonic and ultrasonic activation on organic tissue dissolution

from simulated grooves in root canals using sodium hypochlorite and EDTA. *Int Endod J* 50 (10), 976-982 (2017).

[15] Fariniuk LF, et al: Efficacy of protaper instruments during endodontic retreatment. *Indian J Dent Res* 28 (4), 400-405 (2017).

[16] Giuliani V, Cocchetti R, Pagavino G: Efficacy of ProTaper universal retreatment files in removing filling materials during root canal retreatment. *J Endod* 34 (11), 1381-1384 (2008).

[17] Takahashi CM, et al: In vitro evaluation of the effectiveness of ProTaper universal rotary retreatment system for gutta-percha removal with or without a solvent. *J Endod* 35 (11), 580-3 (2009).