

ZMK 34 (3) 2018, 144-154

Prof. Dr. Claus-Peter Ernst

SDR wird bunt

- [1] Al-Harbi F, Kaisarly D, Bader D, El Gezawi M: Marginal integrity of bulk versus incremental fill Class II composite restorations. *Oper Dent* 41, 146–156 (2016).
- [2] Al Shaafi Haenel T, Sullivan B, Labrie D, Alqahtani MQ, Price RB: Effect of a broad-spectrum LED curing light on the Knoop microhardness of four posterior resin based composites at 2, 4 and 6-mm depths. *J Dent* 45, 14–18 (2016).
- [3] Assis FS, Lima SNL, Tonetto MR, Bhandi SH, Pinto SCS, Malaquias P, Loguercio AD, Bandéca MC: Evaluation of bond strength, marginal integrity, and fracture strength of bulk- vs incrementally-filled restorations. *J Adhes Dent* 18, 317–323 (2016).
- [4] Benetti AR, Havndrup-Pedersen C, Honoré D, Pedersen M, Pallesen U: Bulk-fill resin composites: polymerization contraction, depth of cure, and gap formation. *Oper Dent* 40, 190–200 (2015).
- [5] Bortoluzzi EA, Niu LN, Palani CD et al.: Cytotoxicity and osteogenic potential of silicate calcium cements as potential protective materials for pulpal revascularization. *Dent Mater* 31, 1510–1522 (2015).
- [6] Buchalla W, Lennon AM: Die fluoreszenzunterstützte Kariesexkavation – Funktionsprinzip und Empfehlungen für den Gebrauch. *Oralprophylaxe und Kinderzahnheilkunde* 35, 162–173 (2013).
- [7] Camilleri J, Laurent P, About I: Hydration of biodentine, Theracal LC, and a prototype tricalcium silicate-based dentin replacement material after pulp capping in entire tooth cultures. *J Endod* 40, 1846–1854 (2014).
- [8] Cao Y, Bogen G, Lim J, Shon WJ, Kang MK: Bioceramic materials and the changing concepts in vital pulp therapy. *J Calif Dent Assoc* 44, 278–290 (2016).
- [9] Campos EA, Ardu S, Lefever D, Jassé FF, Bortolotto T, Krejci I: Marginal adaptation of Class II cavities restored with bulk-fill composites. *J Dent* 42, 575–581 (2014).
- [10] Chen C, Niu LN, Xie H, Zhang ZY, Zhou LQ, Jiao K, Chen JH, Pashley DH, Tay FR: Bonding of universal adhesives to dentine – old wine in new bottles? *J Dent* 43, 525–536 (2015).
- [11] Dammaschke T: Dentin- und Hartgewebsneubildung nach Überkappung der Pulpa. *zm* 107 (18), 2084–2094 (2017).
- [12] El-Damanhoury H, Platt J: Polymerization shrinkage stress kinetics and related properties of bulk-fill resin composites. *Oper Dent* 39, 374–382 (2013).
- [13] Ernst CP: Neuversorgung eines insuffizienten Goldinlays. *ZMK* 32, 846–848 (2016).
- [14] Ernst CP: Vom Gold-Standard zur Gold-Alternative. *ZMK* 31, 214–226 (2015).
- [15] Flury S, Hayoz S, Peutzfeldt A, Hüsler J, Lussi A: Depth of cure of resin composites: is the ISO 4049 method suitable for bulk fill materials? *Dent Mater* 28, 521–528 (2012).
- [16] Frankenberger R, Hehn J, Hajto N, Krämer N, Naumann M, Koch A, Roggendorf MJ: Effect of proximal box elevation with resin composite on marginal quality of resin composite inlays in vitro. *Clin Oral Investig* 17, 177–183 (2013).

- [17] Fronza BM, Rueggeberg FA, Braga RR, Mogilevych B, Soares LE, Martin AA, Ambrosano G, Giannini M: Monomer conversion, microhardness, internal marginal adaptation, and shrinkage stress of bulk-fill resin composites. *Dent Mater* 31, 1542–1551 (2015).
- [18] Furness A, Tadros MY, Looney SW, Rueggeberg FA: Effect of bulk/incremental fill on internal gap formation of bulk-fill composites. *J Dent* 42, 439–449 (2014).
- [19] Garcia D, Yaman P, Dennison J, Neiva G: Polymerization shrinkage and depth of cure of bulk fill flowable composite resins. *Oper Dent* 39, 441–448 (2014).
- [20] Garoushi S, Vallittu P, Shinya A, Lassila L: Influence of increment thickness on light transmission, degree of conversion and micro hardness of bulk fill composites. **Quelle?**
- [21] Heintze SD, Monreal D, Peschke A: Marginal quality of class II composite restorations placed in bulk compared to an incremental technique: evaluation with SEM and stereomicroscope. *J Adhes Dent* 17, 147–154 (2015).
- [22] Hirata R, Clozza E, Giannini M, Farrokhmanesh E, Janal M, Tovar N, Coelho PG: Shrinkage assessment of low shrinkage composites using micro-computed tomography. *J Biomed Mater Res B Appl Biomater*, 1–9 (2014).
- [23] Ilgenstein I, Zitzmann NU, Bühler J, Wegehaupt FJ, Attin T, Weiger R, Krastl G: Influence of proximal box elevation on the marginal quality and fracture behavior of root-filled molars restored with CAD/CAM ceramic or composite onlays. *Clin Oral Investig* 19, 1021–1028 (2015).
- [24] Ilie N, Hickel R: Investigations on a methacrylate-based flowable composite based on the SDR technology. *Dent Mater* 27, 348–355 (2011).
- [25] Jang JH, Park SH, Hwang IN: Polymerization shrinkage and depth of cure of bulk-fill resin composites and highly filled flowable resin. *Oper Dent* 40, 172–180 (2015).
- [26] Kalmowicz J, Phebus JG, Owens BM, Johnson WW, King GT: Microleakage of class I and II composite resin restorations using a sonic-resin placement system. *Oper Dent* 40, 653–661 (2015).
- [27] Karaman E, Keskin B, Inan U: Three-year clinical evaluation of class II posterior composite restorations placed with different techniques and flowable composite linings in endodontically treated teeth. *Clin Oral Invest* 21, 709–716 (2017).
- [28] Kapoor N, Bahuguna N, Anand S: Influence of composite insertion technique on gap formation. *J Conserv Dent* 19, 77–81 (2016).
- [29] Kim HJ, Park SH: Measurement of the internal adaptation of resin composites using micro-CT and its correlation with polymerization shrinkage. *Oper Dent* 39, E57–70 (2014).
- [30] Kim RJY, Kim YJ, Choi NS, Lee I-B: Polymerization shrinkage, modulus, and shrinkage stress related to tooth-restoration interfacial debonding in bulk-fill composites. *J Dent* 43, 430–439 (2015).
- [31] Krejci I, Lutz F, Reimer M: Marginal adaptation and fit of adhesive ceramic inlays. *J Dent* 21, 39–46 (1993).
- [32] Kumagai RY, Zeidan LC, Rodrigues JA, Reis AF, Roulet JF: Bond strength of a flowable bulk-fill resin composite in class II MOD cavities. *J Adhes Dent* 17, 427–432 (2015).
- [33] Li X, Pongprueksa P, Van Meerbeek B, De Munck J: Curing profile of bulk-fill resin-based composites. *J Dent* 43, 664–672 (2015).

- [34] Magne P, Spreafico RC: Deep margin elevation: a paradigm shift. *Am J Esthet Dent* 2, 86–96 (2012).
- [35] Manhart, J, Chen H, Hamm, G, Hickel R: Buonocore Memorial Lecture. Review of the clinical survival of direct and indirect restorations in posterior teeth of the permanent dentition. *Oper Dent* 29, 481–508 (2004).
- [36] Manhart J, Chen HY, Hickel R: Clinical evaluation of the posterior composite Quixfil in class I and II cavities: 4-year follow-up of a randomized controlled trial. *J Adhes Dent* 12, 237–243 (2010).
- [37] Marovic D, Tauböck TT, Attin T, Panduric V, Tarle Z: Monomer conversion and shrinkage force kinetics of low-viscosity bulk-fill resin composites. *Acta Odontol Scand* 73, 474–480 (2014).
- [38] Miletic V, Pongprueksa P, De Munck J, Brooks NR, Van Meerbeek B: Curing characteristics of flowable and sculptable bulk-fill composites. *Clin Oral Invest* 21, 1201–1212 (2017).
- [39] Moraschini V, Fai CK, Alto RM, Dos Santos GO: Amalgam and resin composite longevity of posterior restorations: a systematic review and meta-analysis. *J Dent* 43, 1043–1050 (2015).
- [40] Nazari A, Sadr A, Saghiri MA, Campillo-Funollet M, Hamba H, Shimada Y, Tagami J, Sumi Y: Non-destructive characterization of voids in six flowable composites using swept-source optical coherence tomography. *Dent Mater* 29, 278–286 (2013).
- [41] Nazari A, Sadr A, Shimada Y, Tagami J, Sumi Y: 3D assessment of void and gap formation in flowable resin composites using optical coherence tomography. *J Adhes Dent* 15, 237–243 (2013).
- [42] Nowicka A, Lipski M, Parafiniuk M et al.: Response of human dental pulp capped with biodentine and mineral trioxide aggregate. *J Endod* 39, 743–747 (2013).
- [43] Ozel E, Tuna EB, Firatli E: The effects of cavity-filling techniques on microleakage in class II resin restorations prepared with Er:YAG laser and diamond bur: a scanning electron microscopy study. *Scanning* 38, 389–395 (2016).
- [44] Poggio C, Chiesa M, Scribante A, Mekler J, Colombo M: Microleakage in class II composite restorations with margins below the CEJ: In vitro evaluation of different restorative techniques. *Med Oral Patol Oral Cir Bucal* 18, 793–798 (2013).
- [45] Papadogiannis D, Tolidis K, Gerasimou P, Lakes R, Papadogiannis Y: Viscoelastic properties, creep behavior and degree of conversion of bulk fill composite resins. *Dent Mater* 31, 1533–1541 (2015).
- [46] Roggendorf MJ, Krämer N, Appelt A, Naumann M, Frankenberger R: Marginal quality of flowable 4-mm base vs. conventionally layered resin composite. *J Dent* 39, 643–647 (2011).
- [47] Roggendorf MJ, Krämer N, Dippold C, Vosen VE, Naumann M, Jablonski-Momeni A, Frankenberger R: Effect of proximal box elevation with resin composite on marginal quality of resin composite inlays in vitro. *J Dent* 40, 1068–1073 (2012).
- [48] Rosa WL, Piva E, Silva AF: Bond strength of universal adhesives: a systematic review and meta-analysis. *J Dent* 43, 765–776 (2015).
- [49] Rosatto CMP, Bicalho AA, Veríssimo C, Bragança GF, Rodrigues MP, Tantbirojn D, Versluis A, Soares CJ: Mechanical properties, shrinkage stress, cuspal strain and fracture

resistance of molars restored with bulk-fill composites and incremental filling technique. *J Dent* 43, 1519–1528 (2015).

[50] Rullmann I, Schattenberg A, Marx M, Willershausen B, Ernst C-P: Photoelastic determination of polymerization shrinkage stress in low-shrinkage resin composites. *Schweiz Monatsschr Zahnmed* 22, 294–299 (2012).

[51] Rullmann I: Pulpaüberkappung: Untersuchungen zur Wirksamkeit von TheraCal LC. *ZMK* 30 (3), 122–133 (2014).

[52] Schwendicke F, Göstemeyer G, Blunck U, Paris S, Hsu LY, Tu YK: Directly placed restorative materials: review and network meta-analysis. *J Dent Res* 95, 613–622 (2016).

[53] Scotti N, Comba A, Gambino A, Paolino DS, Alovise M, Pasqualini D, Berutti E: Microleakage at enamel and dentin margins with a bulk fills flowable resin. *Eur J Dent* 8, 1–8 (2014).

[54] Scotti N, Alovise C, Comba A, Ventura G, Pasqualini D, Grignolo F, Berutti E: Evaluation of composite adaptation to pulpal chamber floor using optical coherence tomography. *J Endod* 42, 160–163 (2016).

[55] Son S-A, Park J-K, Seo D-G, Ko C-C, Kwon YH: How light attenuation and filler content affect the microhardness and polymerization shrinkage and translucency of bulk-fill composites? *Clin Oral Invest* 21, 559–565 (2017).

[56] Takamizawa T, Barkmeier WW, Tsujimoto A, Scheidel DD, Watanabe H, Erickson RL, Latta MA, Miyazaki M: Influence of water storage on fatigue strength of self-etch adhesives. *J Dent* 43, 1416–1427 (2015).

[57] van Dijken JW, Pallesen U: Posterior bulk-filled resin composite restorations: a 5-year randomized controlled clinical study. *J Dent* 51, 29–35 (2016).

[58] van Dijken JW, Pallesen U: Bulk-filled posterior resin restorations based on stress decreasing resin technology: a randomized, controlled 6-year evaluation. *Eur J Oral Sci* 125, 303–309 (2017).

[59] Van Ende A, De Munck J, Lise DP, Van Meerbeek B: Bulk-fill composites: a review of the current literature. *J Adhes Dent* 9, 95–109 (2017).

[60] Zaruba M, Göhring TN, Wegehaupt FJ, Attin RT: Influence of a proximal margin elevation technique on marginal adaptation of ceramic inlays. *Acta Odontol Scand* 71, 317–324 (2013).

[61] Zorzin J, Maier E, Harre S, Fey T, Belli R, Lohbauer U, Petschelt A, Taschner M: Bulk-fill resin composites: polymerization properties and extended light curing. *Dent Mater* 31, 293–301 (2015).