

ZMK 3 (36) 2020, S 136-146

Prof. Dr. Claus-Peter Ernst

3s-PowerCure-Produktsystem: Hochleistungspolymerisation in 3 Sekunden?

[1] Abbott P, Leow N: Predictable management of cracked teeth with reversible pulpitis. *Aust Dent J* 54, 306–315 (2009).

[2] Al-Harbi F, Kaisarly D, Bader D, El Gezawi M: Marginal integrity of bulk versus incremental fill Class II composite restorations. *Oper Dent* 41, 146–156 (2016).

[3] Assis FS, Lima SNL, Tonetto MR, Bhandi SH, Pinto SCS, Malaquias P, Loguercio AD, Bandéca MC: Evaluation of bond strength, marginal integrity, and fracture strength of bulkvs incrementally-filled restorations. *J Adhes Dent* 18, 317–323 (2016).

[4] Baldissera RA, Corrêa MB, Schuch HS, Collares K, Nascimento GG, Jardim PS, Moraes RR, Opdam NJ, Demarco FF. Are there universal restorative composites for anterior and posterior teeth? *J Dent* 41, 1027–1035 (2013).

[5] Benetti AR, Havndrup-Pedersen C, Honoré D, Pedersen M, Pallesen U: Bulk-fill resin composites: polymerization contraction, depth of cure, and gap formation. *Oper Dent* 40, 190–200 (2015).

[6] Campos EA, Ardu S, Lefever D, Jassé FF, Bortolotto T, Krejci I: Marginal adaptation of Class II cavities restored with bulk-fill composites. *J Dent* 42, 575–581 (2014).

[7] Da Rosa Rodolpho PA, Donassollo TA, Cenci MS, Loguercio AD, Moraes RR, Bronkhorst EM, Opdam NJ, Demarco FF: 22 year clinical evaluation of the performance of two posterior composites with different filler characteristics. *Dent Mater* 27, 955–963 (2011).

[8] El-Damanny H, Platt J: Polymerization shrinkage stress kinetics and related properties of bulk-fill resin composites. *Oper Dent* 39, 374–382 (2013).

[9] Ernst CP: Ersatz eines Goldinlays durch eine Kompositrestauration. *ZMK* 22, 312–316 (2006).

[10] Ernst CP: Kompositrestauration nach Fraktur eines Keramikinlays. *Cosmetic Dentistry* 4 (1), 18–21 (2006).

[11] Ernst CP: Kompositrestauration neben Goldrestauration. *ZMK* 23, 65–67 (2007).

[12] Ernst CP: Komposit im Seitenzahnbereich: Möglichkeiten und Grenzen *Quintessenz* 61, 545–557 (2010).

[13] Ernst CP: Update Komposite – Teil II: Indikation und Anwendung von Kompositmaterialien in der Füllungstherapie. *ZMK* 26, 255–275 (2010).

[14] Ernst CP: Lichtpolymerisation. In: Frankenberger, R.: *Adhäsive Zahnheilkunde*, Deutscher Zahnärzte-Verlag, Köln (2013), 35–80.

[15] Ernst CP: Keramikteilkronen als Versorgungsoption von Zähnen mit Infraktionen. *All*

ceramic partial crowns as a treatment option for cracked, but symptomless teeth. Deutsche Zahnärztl Z 70, 165–173 (2015).

[16] Ernst CP: „Schicht im Schacht“: Aktuelle Betrachtung zum Thema Bulk Fill- versus Schichttechnik bei direkten Seitenzahn-Kompositrestaurationen. ZMK 31, 704–719 (2015).

[17] Ernst CP: Kons. II: Füllungstherapie. Beitrag in: MK Update 2015 – Handbuch ZMK 2015. Herausgeber: Prof. Dr. Grötz, Prof. Dr. Lauer, Prof. Dr. Hellwig, med publico GmbH Wiesbaden, 1–464 (2015).

[18] Ernst CP: Vom Gold-Standard zur Gold-Alternative. ZMK 31, 214–226 (2015).

[19] Ernst CP: Neuversorgung eines insuffizienten Goldinlays. ZMK 32, 846–848 (2016).

[20] Ernst CP: SDR wird bunt. ZMK 34 (3), 144–154 (2018).

[21] Ernst CP: Augen auf beim Lampenkauf. ZMK 35, 14–20 (2019).

[22] Ferracane F, Watts DC, Barghi N, Ernst CP, Rueggeberg FA, Shortall A, Price R; Der effiziente Einsatz von Lichtpolymerisationsgeräten – ein Leitfaden für Zahnärzte. ZMK 30, 166–180 (2014).

[23] Ferracane JL, Hilton TJ: Polymerization stress – Is it clinically meaningful? Dent Mater 32. 1–10 (2016).

[24] Flury S, Hayoz S, Peutzfeldt A, Hüsler J, Lussi A: Depth of cure of resin composites: is the ISO 4049 method suitable for bulk fill materials? Dent Mater 28, 521–528 (2012).

[25] Fronza BM, Rueggeberg FA, Braga RR, Mogilevych B, Soares LES, Martin AA, Ambrosano G, Giannini M: Monomer conversion, microhardness, internal marginal adaptation, and shrinkage stress of bulk-fill resin composites. Dent Mater 31, 1542–1551 (2015).

[26] Furness A, Tadros MY, Looney SW, Rueggeberg FA: Effect of bulk/incremental fill on internal gap formation of bulk-fill composites. J Dent 42, 439–449 (2014).

[27] Gandolfi MG, Siboni F, Prati C: Chemical-physical properties of TheraCal LC, a novel light-curable MTA-like material for pulp capping. International Endodontic Journal 45, 571–579 (2012).

[28] Garcia D, Yaman P, Dennison J, Neiva G: Polymerization shrinkage and depth of cure of bulk fill flowable composite resins. Oper Dent 39, 441–448 (2014).

[29] Garoushi S, Vallittu P, Shinya A, Lassila L: Influence of increment thickness on light transmission, degree of conversion and micro hardness of bulk fill composites. Odontology 104, 291–297 (2016).

[30] Heck K, Manhart J, Hickel R, Diegritz C: Clinical evaluation of the bulk fill composite QuiXfil in molar class I and II cavities: 10-year results of a RCT. Dent Mater 34, e138–e147 (2018).

[31] Heintze SD, Monreal D, Peschke A: Marginal quality of class II composite restorations placed in bulk compared to an incremental technique: evaluation with SEM and

stereomicroscope. *J Adhes Dent* 17, 147–154 (2015).

[32] Hickel R, Ernst CP, Haller B, Hugo B, Kunzelmann KH, Merte K, Ott K, Schmalz G, Staehle HJ: Direkte Kompositrestaurationen im Seitenzahnbereich – Indikation und Lebensdauer. *Zahnärztliche Mitteilungen* 95, 74–76 (2005).

[33] Hickel R, Price R, Soares C, Ilie N: Lichthärtende Bulkfill-Kompositkunststoffe. *zm* 108 (9), 84–85 (2018).

[34] Hirata R, Clozza E, Giannini M, Farrokhmanesh E, Janal M, Tovar N, Coelho PG: Shrinkage assessment of low shrinkage composites using micro-computed tomography. *J Biomed Mater Res B Appl Biomater* 103, 798–806 (2015).

[35] Jacker-Guhr S: Infraktionen. *Dtsch Zahnärztl Z* 69, 429–431 (2014).

[36] Jang JH, Park SH, Hwang IN: Polymerization shrinkage and depth of cure of bulk-fill resin composites and highly filled flowable resin. *Oper Dent* 40, 172–180 (2015).

[37] Kalmowicz J, Phebus JG, Owens BM, Johnson WW, King GT: Microleakage of class I and II composite resin restorations using a sonic-resin placement system. *Oper Dent* 40, 653–661 (2015).

[38] Kapoor N, Bahuguna N, Anand S: Influence of composite insertion technique on gap formation. *J Conserv Dent* 19, 77–81 (2016).

[39] Karaman E, Keskin B, Inan U: Three-year clinical evaluation of class II posterior composite restorations placed with different techniques and flowable composite linings in endodontically treated teeth. *Clin Oral Invest* 21, 709–716 (2017).

[40] Kim HJ, Park SH: Measurement of the internal adaptation of resin composites using micro-CT and its correlation with polymerization shrinkage. *Oper Dent* 39, E57–70 (2014).

[41] Krämer N, Reinelt C, Frankenberger R: Ten-year clinical performance of posterior resin composite restorations. *J Adhes Dent* 17, 433–441 (2015).

[42] Kumagai RY, Zeidan LC, Rodrigues JA, Reis AF, Roulet JF: Bond strength of a flowable bulk-fill resin composite in class II MOD cavities. *J Adhes Dent* 17, 427–432 (2015).

[43] Lempel E, Tóth Á, Fábrián T, Krajczár K, Szalma J: Retrospective evaluation of posterior direct composite restorations: 10-year findings. *Dent Mater* 31, 115–122 (2015).

[44] Li X, Pongprueksa P, Van Meerbeek B, De Munck J: Curing profile of bulk-fill resin-based composites. *J Dent* 43, 664–672 (2015).

[45] Lynch CD, Opdam NJ, Hickel R, Brunton PA, Gurgan S, Kakaboura A, Shearer AC, Vanherle G, Wilson NH: Guidance on posterior resin composites: Academy of Operative Dentistry – European Section. *J Dent* 42, 377–383 (2014).

[46] Magno MB, Rodrigues Nascimento GC, da Rocha YS, Ribeiro BD, Loretto SC, Maia LC: Silorane-based composite resin restorations are not better than conventional composites – a meta-analysis of clinical studies. *J Adhes Dent* 18, 375–386 (2016).

[47] Manhart, J, Chen H, Hamm, G, Hickel R: Buonocore Memorial Lecture. Review of the

clinical survival of direct and indirect restorations in posterior teeth of the permanent dentition. *Oper Dent* 29, 481–508 (2004).

[48] Miletic V, Pongprueksa P, De Munck J, Brooks NR, Van Meerbeek B: Curing characteristics of flowable and sculptable bulk-fill composites. *Clin Oral Invest* 21, 1201–1212 (2017).

[49] Moraschini V, Fai CK, Alto RM, Dos Santos GO: Amalgam and resin composite longevity of posterior restorations: A systematic review and meta-analysis. *J Dent* 43, 1043–1050 (2015).

[50] Nazari A, Sadr A, Saghiri MA, Campillo-Funollet M, Hamba H, Shimada Y, Tagami J, Sumi Y: Non-destructive characterization of voids in six flowable composites using swept-source optical coherence tomography. *Dent Mater* 29, 278–286 (2013).

[51] Nazari A, Sadr A, Shimada Y, Tagami J, Sumi Y: 3D assessment of void and gap formation in flowable resin composites using optical coherence tomography. *J Adhes Dent* 15, 237–243 (2013).

[52] Ozel E, Tuna EB, Firatli E: The effects of cavity-filling techniques on microleakage in class II resin restorations prepared with Er:YAG laser and diamond bur: A scanning electron microscopy study. *Scanning* 38, 389–395 (2016).

[53] Opdam NJ, Roeters JJ, Loomans BA, Bronkhorst EM: Seven-year clinical evaluation of painful cracked teeth restored with a direct composite restoration. *J Endod* 34, 808–811 (2008).

[54] Opdam NJM, van de Sande FH, Bronkhorst E, Cenci MS, Bottenberg P, Pallesen U, Gaengler P, Lindberg A, M.C.D.N.J.M. Huysmans MCDNJM, van Dijken JW: Longevity of posterior composite restorations: A systematic review and meta-analysis. *J Dent Res* 93, 943–949 (2014).

[55] Pallesen U, van Dijken JW: A randomized controlled 30 years follow up of three conventional resin composites in Class II restorations. *Dent Mater* 31, 1232–1244 (2015).

[56] Papadogiannis D, Tolidis K, Gerasimou P, Lakes R, Papadogiannis Y: Viscoelastic properties, creep behavior and degree of conversion of bulk fill composite resins. *Dent Mater* 31, 1533–1541 (2015).

[57] Poggio C, Chiesa M, Scribante A, Mekler J, Colombo M: Microleakage in class II composite restorations with margins below the CEJ: In vitro evaluation of different restorative techniques. *Med Oral Patol Oral Cir Bucal* 18, 793–798 (2013).

[58] Roggendorf MJ, Krämer N, Appelt A, Naumann M, Frankenberger R: Marginal quality of flowable 4-mm base vs. conventionally layered resin composite. *J Dent* 39, 643–647 (2011).

[59] Rosatto CMP, Bicalho AA, Veríssimo C, Bragança GF, Rodrigues MP, Tantbirojn D, Versluis A, Soares CJ: Mechanical properties, shrinkage stress, cuspal strain and fracture resistance of molars restored with bulk-fill composites and incremental filling technique. *J Dent* 43, 1519–1528 (2015).

- [60] Rullmann I, Schattenberg A, Marx M, Willershausen B, Ernst C-P: Photoelastic determination of polymerization shrinkage stress in low-shrinkage resin composites. *Schweiz Monatsschr Zahnmed* 22, 294–299 (2012).
- [61] Rullmann I: Pulpaüberkappung: Untersuchungen zur Wirksamkeit von TheraCal LC. *ZMK* 30 (3), 122–133 (2014).
- [62] Saikaew P, Matsumoto M, Chowdhury A, Carvalho RM, Sano H: Does shortened application time affect long-term bond strength of universal adhesives to dentin? *Oper Dent* 43, 549–558 (2018).
- [63] Sato T, Takagaki T, Ikeda M, Nikaido T, Burrow MF, Tagami J: Effects of selective phosphoric acid etching on enamel using "no-wait" self-etching adhesives. *J Adhes Dent* 20, 407–415 (2018).
- [64] Schwendicke F, Göstemeyer G, Blunck U, Paris S, Hsu LY, Tu YK: Directly placed restorative materials: review and network meta-analysis. *J Dent Res* 95, 613–622 (2016).
- [65] Scotti N, Comba A, Gambino A, Paolino DS, Alovise M, Pasqualini D, Berutti E: Microleakage at enamel and dentin margins with a bulk fills flowable resin. *Eur J Dent* 8, 1–8 (2014).
- [66] Scotti N, Alovise C, Comba A, Ventura G, Pasqualini D, Grignolo F, Berutti E: Evaluation of composite adaptation to pulpal chamber floor using optical coherence tomography. *J Endod* 42, 160–163 (2016).
- [67] Son S-A, Park J-K, Seo D-G, Ko C-C, Kwon YH: How light attenuation and filler content affect the microhardness and polymerization shrinkage and translucency of bulk-fill composites? *Clin Oral Invest* 21, 559–565 (2017).
- [68] van Dijken JW, Pallesen U: A Randomized 10-year prospective follow-up of class II nanohybrid and conventional hybrid resin composite restorations. *J Adhes Dent* 16, 585–592 (2014).
- [69] van Dijken JW, Lindberg A: A 15-year randomized controlled study of a reduced shrinkage stress resin composite. *Dent Mater* 31, 1150–1158 (2015).
- [70] van Dijken JW, Pallesen U: Posterior bulk-filled resin composite restorations: A 5-year randomized controlled clinical study. *J Dent* 51, 29–35 (2016).
- [71] van Dijken JW, Pallesen U: Bulk-filled posterior resin restorations based on stress decreasing resin technology: a randomized, controlled 6-year evaluation. *Eur J Oral Sci* 125, 303–309 (2017).
- [72] van Ende A, De Munck J, Lise DP, Van Meerbeek B: Bulk-fill composites: a review of the current literature. *J Adhes Dent* 9, 95–109 (2017).
- [79] Zorzini J, Maier E, Harre S, Fey T, Belli R, Lohbauer U, Petschelt A, Taschner M: Bulk-fill resin composites: polymerization properties and extended light curing. *Dent Mater* 31, 293–301 (2015).