

Ästhetik ist eine Korrelation zwischen Form und Funktion – Ein integriertes Konzept für kieferorthopädisch-kieferchirurgische Therapie von skelettalen Dysgnathien

- [1] Albino JE, Tedesco L: Esthetic need for orthodontic treatment. In Melsen B, editor: Current controversies in orthodontics. Chicago, Quintessence Publishing, 11–24 (1994).
- [2] Al-Riyami S, Cunningham SJ, Moles DR: Orthognathic treatment and temporomandibular disorders: a systematic review, Part 2. Signs and symptoms and meta-analyses. Am J Orthod Dentofacial Orthop 136 (5), 626.e1–16, discussion 626–627 (2009).
- [3] Bass NM: Dento-facial orthopaedics in the correction of the skeletal II malocclusion. Br J Orthod 9, 3–8 (1982).
- [4] Bell HW, Jacobs JD: Simultaneous repositioning of the maxilla, mandible, and chin. Treatment planning and analysis of the soft tissues. Am J Orthod 91, 28–50 (1986).
- [5] Bell WH, Proffit HR, White RP: Surgical correction of facial deformities. Saunder Verlag, Baltimor (1980).
- [6] Bell WH, Schdeman GB: Correction of vertical maxillary deficiency: Stability and soft tissue changes. J Oral Surg 39, 666–670 (1981).
- [7] Berscheid E, Gangestade S: The social psychological implications of facial physical attractiveness. Clin Plast Surg 9, 289–296 (1982).
- [8] Buddgaard M, Melsen B, Terp S: Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. Eur J Orthod 8, 21–29 (1986).
- [9] Canut J: Eine Analyse der dentofazialen Ästhetik. Inf Orthod Kieferorthop 28, 83–105 (1996).
- [10] Centenero SAH, Hernández-Alfaro F: 3D planning in orthognathic surgery: CAD/CAM surgical splints and prediction of the soft and hard tissues results – Our experience in 16 cases. J Craniomaxillofac Surg 40, 162–168 (2012).
- [11] Collins P, Epker BN: The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 53, 549–554 (1982).
- [12] Dal Pont G: L'osteotomia retromolare per la correzione della prognathia, Minerva chir 18, 1138–1141 (1959).
- [13] Dal Pont G: Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 58, 8–10 (1961).
- [14] Dryland-Vig KWL, Ellis III E: Diagnosis and treatment planning for the surgical-orthodontic Patient. Cli Plast Surg 16, 645–658 (1989).
- [15] Dann JJ, Fonseca RJ, Bell WH: Soft tissue changes associated with total maxillary advancement: a preliminary study. J Oral Surg 34, 19–23 (1976).
- [16] De Assis EA, Starck WJ, Epker B: Cephalometric analysis of profile nasal esthetics. Part III. Postoperativ changes after isolated superior repositioning. Int J Adult Orthod Orthognath 11, 279–288 (1996).
- [17] Ellis E: Condylar positioning devices for orthodontic surgery: Are they necessary. J Oral Maxillofac Syrg 52, 536–552 (1994).
- [18] Ellis E, Hinton RJ: Histologic examination of the temporomandibular joint after mandibular advancement with and without rigid fixation: An experimental investigation in adult *Maccaca mulatta*. J Oral Maxillofac Syrg 49, 1316 (1991).
- [19] Epker B: Esthetic maxillofacial surgery. Lea und Febiger Verlag, Philadelphia (1994).
- [20] Epker BN, Fish LC: Surgical-orthodontic correction of open-bite deformity. Am J Orthod 71, 278–299 (1977).
- [21] Epker BN, Fish LC: Surgical superior repositioning of the maxilla: What to do with the mandible? Am J Orthod 78, 164–191 (1980).

- [22] Farkas LG, Kolar JC: Anthropometry and art in the aesthetics of women's face. *Clin Plast Surg* 14, 599–615 (1987).
- [23] Flanary CM, Barnwell GM, Alexander JM: Patient perceptions of orthognathic surgery. *Am J Orthod* 88, 137–145 (1985).
- [24] Freihofer HPM: Changes in nasal profile after maxillary advancement in cleft and non-cleft patients. *J Maxillofac Surg* 5, 20–27 (1977).
- [25] Helm S, Siersbaek-Nielsen S, Skieller V, Björk A: Skelatal maturation of the hand in relation to maximum puberal growth in body height. *Danish Dental Journal* 75, 1223–1234 (1971).
- [26] Hogeman KE, Sarnäs KV, Willmar K: One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. *Fortschr Kiefer Gesichtschr* 18, 39–49 (1974).
- [27] Hui E, Hägg EUO, Tidman H: Soft tissue changes following maxillary osteotomies in cleft lip and palate and non-cleft Patients. *J Craniomaxillofac Surg* 22, 182–186 (1994).
- [28] Jensen AC, Sinclair PM, Wolford LM: Soft tissue changes associated with double jaw surgery. *Am J Orthod Dentofac Orthop* 101, 266–275 (1992).
- [29] Jacobson A: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod* 79, 399–415 (1981).
- [30] Joss CU, Vassalli IM: Stability after bilateral sagittal split osteotomy advancement surgery with rigid internal fixation: a systematic review. *J Oral Maxillofac Surg* 67, 301–313 (2009).
- [31] Kiyak HA, Hohl T, West RA: Psychologie changes in orthognathic surgery patients: a 24-month follow-up. *J Oral Maxillofac Surg* 42, 506–512 (1984).
- [32] Lazaridou-Terzoudi T, Kiyak HA, Moore R, Athanasiou AE, Melsen B: Long-term assessment of psychologic outcomes of orthognathic surgery. *J Oral Maxillofac Surg* 61, 545–552 (2003).
- [33] Legan HL, Burstone GJ: Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 38, 744–751 (1980).
- [34] Lee DY, Bailey LJ, Proffit WR: Soft tissue changes after superior repositioning of the maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 11, 301–311 (1996)-
- [35] McNamara JA, McDougall Jr PD, Dierks JM: Arch with development in Class II patients treated with extraoral force and functional jaw orthodontics. *Am J Orthodont* 52, 353–359 (1966).
- [36] Mensink G, Verweij JP, Frank MD, Eelco BJ, Richard van Merkesteyn JP: Bad split during bilateral sagittal split osteotomy of the mandible with separators: a retrospective study of 427 patients. *Br J Oral Maxillofac Surg* 51, 525–529 (2013).
- [37] Mensink G, Zweers A, Wolterbeek R, Dicker GG, Groot RH, van Merkesteyn RJ: Neurosensory disturbances one year after bilateral sagittal split osteotomy of the mandibula performed with separators: A multi-centre prospective study. *J Craniomaxillofac Surg* 40, 763–767 (2012).
- [38] Obwegeser H: The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. *J Oral Surg* 10, 677–689 (1957).
- [39] Obwegeser H: The indication for surgical correction of mandibular deformity by sagittal splitting technique. *Br J Surg* 1, 157–168 (1963).
- [40] Obwegeser H, Trauner R: Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. *Dtsch Zahn Mund Kieferheilk* 23, 1–26 (1955).
- [41] Petrovic AG, Stutzmann J: Reaktionsfähigkeit des tierischen und menschlichen Kondylenknorpels auf Zell- und Molekularebene im Lichte einer kybernetischen Auffassung des faszialen Wachstums. *Fortschr Kieferorthop* 49, 405–425 (1988).
- [42] Proothi M, Drew SJ, Sachs SA: Motivating factors for patients undergoing orthognathic surgery evaluation. *J Oral Maxillofac Surg* 68, 1555–1559 (2010).
- [43] Radney LJ, Jacobs JD: Soft tissue changes associated with surgical total maxillary intrusion. *Am J Orthodont* 80, 191–212 (1981).

- [44] Rosen HM: Lip-nasal esthetics following Le Fort I osteotomy. *Plast Reconstr Surg* 81, 171–182 (1988).
- [45] Schwarz AM: Die Röntgendiagnostik. Urban & Schwarzenberg, Wien (1958).
- [46] Scott O, Kijak HA: Treatment expectation versus outcomes among orthodontic surgery patients. *Int J Adult Orthod Orthognath Surg* 6, 247–255 (1991).
- [47] Solano-Hernandez B, Antonarakis GS, Scolozzi P, Kiliaridis S: Combined orthodontic and orthognathic surgical treatment for the correction of skeletal anterior open-bite malocclusion: a systematic review on vertical stability. *J Oral Maxillofac Surg* 71 (1), 98–109 (2013).
- [48] Watted N, Witt E: NMR study of TNJ changes following functional orthopaedic treatment using the „Würzburg approach“, European Orthodontic Society (EOS) 74th Congress (1998).
- [49] Watted N: Behandlung von Klasse-II-Dysgnathien – funktionskieferorthopädische Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik. *Kieferorthop* 13, 193–208 (1999).
- [50] Watted N, Bill J, Witt E: Therapy concept for the combined orthodontic-surgical treatment of Angle class II deformities with short face syndrome. New aspects for surgical lengthening of the lower face. *Clin Orthod Res* 3, 78–93 (2000).
- [51] Watted N, Bill J, Witt E, Reuther J: Lengthening of the lower face Angle class II patients with skeletally deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbourg, France(1999).
- [52] Watted N, Teuscher T, Wieber M: Vertikaler Gesichtsaufbau und Planung kieferorthopädisch-kieferchirurgischer Kombinationsbehandlungen unter besonderer Berücksichtigung der dentofazialen Ästhetik. *Kieferorthop* 16, 29–44 (2000).
- [53] Westermark AH, Bystedt H, von Konow L, Sällström KO: Nasolabial morphology after Le Fort I osteotomies. *Int J Oral Maxillofac Surg* 20, 25–30 (1991).
- [54] Williamsone EH, Caves SA, Edenfield RJ, Morse PK: Cephalometric analysis: comparisons between maximum intercuspal and centric relation. *Am J Orthod* 74, 672–677 (1978).
- [55] Williamsone EH, Evans DL, Barton WA, Williams BH: The effect of bite plane use on terminal hinge axis location. *Angle Orthod* 47, 25–33 (1977).
- [56] Witt E: Möglichkeiten und Grenzen der kieferorthopädischen Behandlung Erwachsener. *Fortschr Kieferorthop* 52, 1–7 (1991).
- [57] Witt E: Behandlungskonzepte. In Miethke RR, Drescher D (Hrsg.): Kleines Lehrbuch der Angel-Klasse II/1 unter besonderer Berücksichtigung der Behandlung. Quintessenz, Berlin (1996).