

Wirksamkeit noninvasiver Therapien zur Inaktivierung oder Reduktion der Wurzelkaries

- [1] Baca P, et al: Effect of chlorhexidine-thymol varnish on root caries in a geriatric population: a randomized double-blind clinical trial. *J Dent* 37 (9), 679–85 (2009).
- [2] Banoczy J, Nemes J: Effect of amine fluoride (AmF)/stannous fluoride (SnF₂) toothpaste and mouthwashes on dental plaque accumulation, gingivitis and root-surface caries. *Proc Finn Dent Soc* 87 (4), 555–9 (1991).
- [3] Banting DW, et al: The effectiveness of 10 % chlorhexidine varnish treatment on dental caries incidence in adults with dry mouth. *Gerodontology* 17 (2), 67–76 (2000).
- [4] Baysan A, Lynch E: Clinical reversal of root caries using ozone: 6-month results. *Am J Dent* 20 (4), 203–8 (2007).
- [5] Baysan A, et al: Reversal of primary root caries using dentifrices containing 5,000 and 1,100 ppm fluoride. *Caries Res* 35 (1), 41–6 (2001).
- [6] Davey Smith G, Egger M: Meta-analysis. Unresolved issues and future developments. *BMJ* 316 (7126), 221–5 (1998).
- [7] DePaola PF: Caries in our aging population: What are we learning?, in *Cariology for the nineties*, Bowden GH and Tabak LA, Editors. University of Rochester Press: Rochester, NY. 25–35 (1993).
- [8] Ekstrand KR, et al: A Randomized Clinical Trial of the Anti-Caries Efficacy of 5,000 Compared to 1,450 ppm Fluoridated Toothpaste on Root Caries Lesions in Elderly Disabled Nursing Home Residents. *Caries Res* 47 (5), 391–398 (2013).
- [9] Ettinger RL: Epidemiology of dental caries. A broad review. *Dent Clin North Am* 43 (4), 679–94 (1999).
- [10] Fure S, Gahnberg L, Birkhed D: A comparison of four home-care fluoride programs on the caries incidence in the elderly. *Gerodontology* 15 (2), 51–60 (1998).
- [11] Fure S, Lingstrom P: Evaluation of different fluoride treatments of initial root carious lesions in vivo. *Oral Health Prev Dent* 7 (2), 147–54 (2009).
- [12] Geiger AM, et al: Reducing white spot lesions in orthodontic populations with fluoride rinsing. *Am J Orthod Dentofacial Orthop* 101 (5), 403–7 (1992).
- [13] Griffin SO, et al: Estimating Rates of New Root Caries in Older Adults. *J Dent Res* 83 (8), 634–638 (2004).
- [14] Guyatt GH, et al: GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336 (7650), 924–6 (2008).
- [15] Higgins JPT, Green SG: *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5. 10 [updated March 2011]. The Cochrane Collaboration. Available from www.cochrane-handbook.org. 2011.
- [16] Holmes J: Clinical reversal of root caries using ozone, double-blind, randomised, controlled 18-month trial. *Gerodontology* 20 (2), 106–14 (2003).
- [17] Hu DY, et al: A clinical investigation of the efficacy of a dentifrice containing 1.5 % arginine and 1450 ppm fluoride, as sodium monofluorophosphate in a calcium base, on primary root caries. *J Clin Dent* 24 Spec no A, A23–31 (2013).
- [18] Jensen ME, Kohout F: The effect of a fluoridated dentifrice on root and coronal caries in an older adult population. *J Am Dent Assoc* 117 (7), 829–32 (1988).

- [19] Marinho VC, et al: Topical fluoride (toothpastes, mouthrinses, gels or varnishes) for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev* (4), CD002782 (2003).
- [20] Michaelis W, Schiffner U: *The Fourth German Oral Health Study (DMS IV)*. Institute of German Dentists (IDZ), Deutscher Zahnärzte Verlag: Köln (2006).
- [21] Nitschke I: Zahnmedizinische Grundlagen zur geriatrischen Rehabilitation – eine Einführung in die Alternszahnmedizin. *Zeitschrift für Gerontologie und Geriatrie* 33 (1 Supplement), 45–049 (2000).
- [22] Papas A, et al: Comparative efficacy of stabilized stannous fluoride/sodium hexametaphosphate dentifrice and sodium fluoride/triclosan/copolymer dentifrice for the prevention of periodontitis in xerostomic patients: a 2-year randomized clinical trial. *J Periodontol* 78 (8), 1505–14 (2007).
- [23] Papas A, et al: Caries clinical trial of a remineralising toothpaste in radiation patients. *Gerodontology* 25 (2), 76–88 (2008).
- [24] Paraskevas S, et al: Amine fluoride/stannous fluoride and incidence of root caries in periodontal maintenance patients. A 2 year evaluation. *J Clin Periodontol* 31 (11), 965–71 (2004).
- [25] Peters MC,: Strategies for noninvasive demineralized tissue repair. *Dent Clin North Am* 54 (3), 507–25 (2010).
- [26] Petersson LG, et al: Remineralization of primary root caries lesions using an amine fluoride rinse and dentifrice twice a day. *Am J Dent* 20 (2), 93-6 (2007).
- [27] Powell LV, et al: Caries prevention in a community-dwelling older population. *Caries Res* 33 (5), 333–9 (1999).
- [28] Ravald N, Birkhed D: Prediction of root caries in periodontally treated patients maintained with different fluoride programmes. *Caries Res* 26 (6), 450–8 (1992).
- [29] Ripa LW, et al: Effect of a 0.05 % neutral NaF mouthrinse on coronal and root caries of adults. *Gerodontology* 6 (4), 131–6 (1987).
- [30] Schaeken MJ, Keltjens HM, Van Der Hoeven JS: Effects of fluoride and chlorhexidine on the microflora of dental root surfaces and progression of root-surface caries. *J Dent Res* 70 (2), 150–3 (1991).
- [31] Shellis RP: A scanning electron-microscopic study of solubility variations in human enamel and dentine. *Arch Oral Biol* 41 (5), 473–84 (1996).
- [32] Souza ML, et al: Comparing the efficacy of a dentifrice containing 1.5 % arginine and 1450 ppm fluoride to a dentifrice containing 1450 ppm fluoride alone in the management of primary root caries. *J Dent* 41 Suppl 2, S35–41 (2013).
- [33] Srinivasan M, et al: High-fluoride toothpaste: a multicenter randomized controlled trial in adults. *Community Dent Oral Epidemiol* (2013).
- [34] Tan HP, et al: A randomized trial on root caries prevention in elders. *J Dent Res* 89 (10), 1086–90 (2010).
- [35] US Department of Health and Human Services, Oral Health in America: A Report of the Surgeon General. Rockville, MD: US Department of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health, 308 (2000).
- [36] Vered Y, et al: Comparison of a dentifrice containing 0.243 % sodium fluoride, 0.3 % triclosan, and 2.0 % copolymer in a silica base, and a dentifrice containing 0.243 % sodium fluoride in a silica base: a three-year clinical trial of root caries and dental crowns among adults. *J Clin Dent* 20 (2). 62–5 (2009).

- [37] Wallace MC, Retief DH, Bradley EL: The 48-month increment of root caries in an urban population of older adults participating in a preventive dental program. *J Public Health Dent* 53 (3), 133–7 (1993).
- [38] Wierichs RJ, Meyer-Lueckel H: Systematic review on noninvasive treatment of root caries lesions. *J Dent Res* 94 (2), 261–71 (2015).
- [39] Wyatt CC, MacEntee MI: Caries management for institutionalized elders using fluoride and chlorhexidine mouthrinses. *Community Dent Oral Epidemiol* 32 (5), 322–8 (2004).
- [40] Wyatt CC, et al: Chlorhexidine and preservation of sound tooth structure in older adults. A placebo-controlled trial. *Caries Res* 41 (2), 93–101 (2007).
- [41] Zhang W, et al: Silver Diamine Fluoride and Education to Prevent and Arrest Root Caries among Community-Dwelling Elders. *Caries Res* 47 (4), 284–90 (2013).